

# ***Bacteroides*: From a Fecal Pollutant to a Useful Tool in Solving Water Pollution**

Christian Jordan O. Dela Rosa and Windell L. Rivera\*

Pathogen-Host-Environment Interactions Research Laboratory, Institute of Biology, College of Science, University of the Philippines Diliman, Quezon City 1101, Philippines

## ABSTRACT

**B***acteroides* spp. are Gram-negative bacilli, non-endospore-forming and obligate anaerobes, belonging to Phylum Bacteroidetes, Class Bacteroidia, Order Bacteroidales, Family *Bacteroidaceae*, with 40 taxonomically recognized species. They are anaerobes, and hence, commonly found in warm-blooded animals' guts and fecal microbiome, where they are the most dominant bacteria, having either mutually or commensally co-evolved with their host. However, they are also disease-causing and carriers of antibiotic resistance genes. Hence, release of *Bacteroides*-dominated feces by hosts causes fecal pollution, especially in water bodies. Despite being anaerobes, *Bacteroides* have a short lifespan. They leave their genetic materials behind as environmental DNA, which remain viable for PCR. DNA isolation can determine the feces' source—a technique known as microbial source tracking (MST). There are two types of MST: library-dependent (LDM) and -independent (LIM). *Bacteroides* use falls under LIM, wherein animal associated *Bacteroides* markers determine the fecal contamination's source.

Due to extensive co-evolution between *Bacteroides* and hosts, most LIM-MST biomarkers are *Bacteroides*-based [*PigBac* (pigs), *CowBac* (cows), *ChickenBac* (chicken)] and human-based (*HF183*, *HumBac*), etc. MST also effectively determines fecal pollution extent from anthropogenic and agricultural sources. Therefore, MST can directly determine a definitive fecal pollutant source. The main limitation of LIM-MST is the lack of markers for other animals. Most available MST markers are directed toward domesticated animals (chickens, cows, pigs). Moreover, even existing markers are affected by geography and animal diet. These limitations warrant continued efforts to fill the gaps by designing more specific and sensitive MST animal markers. This review aims to initiate interest in the use of *Bacteroides* for efficient water quality monitoring.

## INTRODUCTION

The gut contains a rich microbial ecosystem that plays a key role in animal health (Turroni et al. 2020; Zafar and Saier 2021). The human intestine alone contains approximately  $10^{11}$  to  $10^{12}$  bacteria per gram of colonic content (Eckburg et al. 2005).

\*Corresponding author

Email Address: wlriviera@science.upd.edu.ph

Date received: September 24, 2023

Date revised: November 6, 2023

Date accepted: November 21, 2023

## KEYWORDS

*Bacteroides*, fecal pollution, microbial source tracking (MST), library-independent MST, MST markers

Numerically, the most dominant group of bacteria in the human intestine and other warm-blooded animals belongs to the Order Bacteroidales (Kabiri et al. 2013; Coyne et al. 2014; Tamana et al. 2021; Zafar and Saier 2021). *Bacteroides* exclusively live and grow in their animal host's gut, suggesting a strong evolutionary adaptation to the gastrointestinal tract (GIT) environment (Wexler and Goodman 2017). *Bacteroides* are introduced into its human host via vaginal or cesarian delivery and skin contact with lactating mother (Wang et al. 2015; Milani et al. 2017). *B. thetaiotaomicron* is commonly found to be the prevalent *Bacteroides* during the lactating months of a human baby since this *Bacteroides* species can highly degrade glycan from human milk (Fouhy et al. 2012; Milani et al. 2017). As more solid foods are introduced, more *Bacteroides* species inhabit the niches in the gut since they are highly specialized in the degradation of several kinds of glycan. As humans grow and consume more food groups, particularly during childhood, *Bacteroides* species shift dominance to other species such as *B. ovatus* and *B. xylanisolvens* (Zhong et al. 2019). Differences in diets among adults show differences in dominant *Bacteroides* species. Among omnivorous adult humans, *B. salanitronis* and *B. coprocola* co-dominate in the gut. Conversely, for vegetarian adults, *B. vulgatus* is dominant while *B. salyersiae* is prevalent in vegans (Ferrocino et al. 2015). However, *B. thetaiotaomicron* and *B. fragilis* are generally found in high concentrations in all stages of life (Zafar and Saier 2021).

*Bacteroides* in the gut play beneficial roles in their warm-blooded animal hosts, although *Bacteroides* have already been isolated from cold-blooded animals such as fishes and frogs (Kabiri et al. 2013; Gibb et al. 2017). They have been recognized either as mutualists or commensals by providing nutritional and immunological benefits, thus establishing stable long-term beneficial relationships with their hosts. Thus, *Bacteroides* are considered clinically important anaerobes that are directly related to intestinal well-being (Hong et al. 2008). They are so extensive and highly adapted to life and survival in the gut that they have evolutionarily become an essential species in the gut microbial food web (Wexler and Goodman 2017). However, they can also cause harm like abscesses to infected organs and bacteremia when they break out of the gut. Moreover, *Bacteroides* undergo extensive horizontal gene transfer (HGT) that include virulence genes and antimicrobial resistomes. Resistomes are well characterized collection of antibiotic resistance genes (ARGs) from a variety of environments (Ma et al. 2021). Some of the ARGs are transferred to other *Bacteroides* spp. or other transient bacteria of the gut, and thus, when these *Bacteroides* or the transient bacteria are expelled with the feces, they have with them the antibiotic resistance phenotype

(Huddleston 2014). *Bacteroides* constitute 30%–50% of the total fecal bacteria, with concentrations of  $10^{11}$  cells per gram of feces (Paruch et al. 2019). Although only a small portion of the gut microbial population is culturable, 14%–40% of cultivable microorganisms in the fecal microbiota are *Bacteroides* (Hong et al. 2008). Due to the dominance of *Bacteroides* in both the gut and fecal microbiome and its strong evolutionary ties to its host, it has been used as an alternative fecal bacterium for monitoring water pollution, instead of the traditional fecal indicator bacteria (FIB) such as the fecal coliforms (*Escherichia coli* and *Enterococcus faecalis*). Hence, this review discusses *Bacteroides* as a fecal pollutant that contributes to the degradation of water quality and causes illnesses, to a more positive role of being part of the solution by identifying the fecal origin – a step forward for developing better governmental policies in addressing the problem of water pollution.

### Taxonomy and biology of *Bacteroides*

Taxonomically, *Bacteroides* sp. belongs to Phylum Bacteroidetes, Class Bacteroidia, Order Bacteroidales, and Family Bacteroidaceae. In a review by Wexler (2007), there were 21 taxonomically known species of *Bacteroides*. Those were *B. acidifaciens*, *B. caccae*, *B. coprocola*, *B. coprosuis*, *B. eggerthii*, *B. finegoldii*, *B. fragilis*, *B. helcogenes*, *B. intestinalis*, *B. massiliensis*, *B. nordii*, *B. ovatus*, *B. thetaiotaomicron*, *B. vulgatus*, *B. plebius*, *B. uniformis*, *B. salyersae*, *B. pyogenes*, *B. goldstenii*, *B. dorei*, and *B. johnsonii*. Through the succeeding years, the number of *Bacteroides* species reached 136 (<https://lpsn.dsmz.de/>). However, further research on each *Bacteroides* shows that out of 136, only 40 are taxonomically legitimate as enumerated in Table 1. The remaining 96 “*Bacteroides*” were either taxonomically transferred to a different genus and have simply become a taxonomic synonym or were not validly published. For example, *B. massiliensis* has been changed to *Phocaeicola massiliensis*, *B. goldsteineii* to *Parabacteroides goldsteini*, and *B. dorei* to *Phocaeicola dorei*. Moreover, some that were identified as “*Bacteroides*” are not considered taxonomically legitimate, yet they were not validly published. This includes *B. congoensis*, *B. cutis*, and *B. sediment*, etc.

Although *Bacteroides* is the most predominant bacterial group in the gut, there are only 40 taxonomically legitimate known species compared with other bacteria. This may be because *Bacteroides* are anaerobes, and anaerobic conditions are one of the most challenging conditions to be able to obtain an enriched or pure culture (Mori and Kamagata 2014).

**Table 1: List of 40 *Bacteroides* valid species**

<i>Bacteroides</i> species	GenBank Accession Number	Source	Reference
1. <i>B. acidifaciens</i>	AB510696	Mice cecum	Miyamoto and Itoh 2000
2. <i>B. caccae</i>	X83951	Human feces	Johnson et al. 1986
3. <i>B. caecicola</i>	AB910337	Chicken cecum	Irisawa et al. 2016
4. <i>B. caecigallinarum</i>	AB861981	Chicken cecum	Saputra et al. 2015
5. <i>B. caecimuris</i>	KR364741	Mice intestine	Lagkouvardos et al. 2016
6. <i>B. cellulolyticus</i>	AJ583243	Human gut	Robert et al. 2007
7. <i>B. clarus</i>	AB490801	Human feces	Watanabe et al. 2010
8. <i>B. coprosuis</i>	AB510699	Pig manure	Whitehead et al. 2005
9. <i>B. eggerthii</i>	AB050107	Human feces	Holdeman and Moore 1974 (Approved Lists 1980)
10. <i>B. faecalis</i>	MK207058	Human feces	Yu et al. 2019

11.	<i>B. faecichinchillae</i>	AB574480	Chinchilla feces	Kitahara et al. 2012
12.	<i>B. faecis</i>	AB547640	Human feces	Kim et al. 2010
13.	<i>B. finegoldii</i>	AB222699	Human feces	Bakir et al. 2006a
14.	<i>B. fluxus</i>	AB490802	Human feces	Watanabe et al. 2010
15.	<i>B. fragilis</i>	AB050106	Human intestine	Castellani and Chalmers 1919 (Approved Lists 1980)
16.	<i>B. galacturonicus</i>	DQ497994	Human intestine	Jensen and Canale-Parola 1986
17.	<i>B. gallinaceum</i>	AB910339	Chicken cecum	Irisawa et al. 2016
18.	<i>B. gallinarum</i>	AB253732	Chicken cecum	Lan et al. 2006
19.	<i>B. graminisolvens</i>	AB363973	Cow waste	Nishiyama et al. 2009
20.	<i>B. helcogenes</i>	AB200227	Pig feces	Benno et al. 1983
21.	<i>B. intestinalis</i>	AB214328	Human feces	Bakir et al. 2006b
22.	<i>B. koreensis</i>	KX025133	Human feces	Shin et al. 2017
23.	<i>B. kribbi</i>	KX025134	Human feces	Shin et al. 2017
24.	<i>B. luhongzhouii</i>	MK584158	Human feces	Ge et al. 2021
25.	<i>B. luti</i>	AB787271	Anaerobic treatment sludge/ facility	Hatamoto et al. 2014
26.	<i>B. nordii</i>	AB510704	Human intestine	Song et al. 2004
27.	<i>B. oleiciplenus</i>	AB490803	Human feces	Watanabe et al. 2010
28.	<i>B. ovatus</i>	X83952	Human feces	Eggerth and Gagnon 1932 (Approved Lists 1980)
29.	<i>B. pectinophilus</i>	ABVQ01000036	Human intestine	Jensen and Canale-Parola 1986
30.	<i>B. propionicifaciens</i>	AB264625	Cow feces	Ueki et al. 2008
31.	<i>B. pyogenes</i>	AB200229	Pig feces	Benno et al. 1983
32.	<i>B. reticulotermis</i>	AB692943	Termite gut	Sakamoto and Ohkuma 2013
33.	<i>B. rodentium</i>	AB547646	Chinchilla feces	Kitahara et al. 2011
34.	<i>B. salyersiae</i>	AB510707	Human intestine	Song et al. 2004
35.	<i>B. stercorisoris</i>	AB574481	Chinchilla feces	Kitahara et al. 2012
36.	<i>B. stercoris</i>	X83953	Human feces	Johnson et al. 1986
37.	<i>B. thetaiotaomicron</i>	AB050109	Human gut	Castellani and Chalmers 1919 (Approved Lists 1980)
38.	<i>B. uniformis</i>	AB050110	Human feces	Eggerth and Gagnon 1932 (Approved Lists 1980)
39.	<i>B. xylanisolvens</i>	AB510713	Human feces	Chassard et al. 2008
40.	<i>B. zhangwenhongii</i>	CP059856	Human feces	Ge et al. 2021

Morphologically, *Bacteroides* are Gram-negative, non-motile, rod-shaped, and non-spore-forming bacteria, as consistently shown in all references enumerated in Table 1. Based on these references, most pure culture isolations were performed using Eggerth-Gagnon (EG) Agar supplemented with 5% blood. However, several recent publications on *Bacteroides* isolation use *Bacteroides* Bile Esculin (BBE) agar (Niestepski et al. 2019; Zamani et al. 2020; Dela Rosa and Rivera 2021). Interestingly, the colonial morphology is circular, entire, raised convex, smooth, and grayish to off-white for both EG and BBE agars. In all instances, the culture condition is at 37°C for 48–72 h in an anaerobic condition to simulate the submicromolar range of oxygen levels in the large intestine. Besides its anaerobic physiology, due to the challenging environment of the large

intestine, where most of the gut microorganisms' live, *Bacteroides* have developed other adaptive mechanisms for their survival in the adverse environment of the large intestine. For example, many species of *Bacteroides* contain cytochrome *bd* oxidase that uses oxygen as a terminal electron acceptor, thus, lowering oxygen levels in the gut. Interestingly, some *Bacteroides* contain the gene *nrdA*, which codes for an enzyme used only in aerobic respiration. Through these, *Bacteroides* can tolerate oxygen and allow their spread to new hosts. (Smalley et al. 2002; Wexler and Goodman 2017).

## **Bacteroides and the vertebrate gut**

### *Bacteroides as mutualist and commensal*

Intestinal *Bacteroides* have co-evolved intricate symbiotic relationships with their specific animal hosts (Bäckhed et al. 2005). The interaction is characterized by *Bacteroides* producing secondary metabolites essential to its host, and in return, the animal host providing stable gut microbial ecosystem (Groussin et al. 2021). Moreover, to a limited extent, *Bacteroides* are also involved in developing the host's immune system (Troy and Kasper 2010). In comparing the fecal microbiota of different host vertebrates using 16S rRNA gene, it showed that *Bacteroides* spp. co-evolve with their hosts, and co-diversified with them (Ley et al. 2008). Symbiosis is so strong that there is a development of host-microbiome fidelity and cohesive association (Groussin et al. 2017).

One of the well-researched symbiotic relationships between *Bacteroides* and its human host is glycans – simple and complex carbohydrates that play metabolic, structural, and physical role in biological systems (Comstock 2009; Varki 2017). The mammalian host lacks the enzymatic capacity to degrade glycans, however, *B. thetaiotamicron*, a glycopile, can break down various dietary glycans. It has 88 polysaccharide utilization loci for degrading various glycans, including plant glycans/polysaccharides (Guo et al. 2020). This capability is not exclusive to *B. thetaiotamicron*. The collective *Bacteroides* population poses an enormous, combined polysaccharide degradative ability capable of metabolizing more than a dozen different kinds of plant and host-derived polysaccharides (Comstock 2009; Wexler and Goodman 2017).

More recently, *B. thetaiotamicron* was proven to restrict the growth of *Clostridioides difficile* colonization in the gut. There is a direct association that *B. thetaiotamicron* increases the concentration of several acids ( $\alpha$ -muricholic,  $\beta$ -muricholic, ketolithocholic, and deoxycholic) known to inhibit *C. difficile* and reduces the taurocholic acid production that promotes *C. difficile* germination (Li et al. 2021b).

On the other hand, *B. fragilis*, conversely, exerts a beneficial immunologic effect. In an experiment conducted on germ-free gut mice, it was discovered that the mice have smaller Payer's patches, fewer germinal centers, reduced IgA-producing plasma cells, decreased CD4<sup>+</sup> T cells in the intestinal lamina propria, and structural defects of the splenic and lymphoid nodes. However, after orally introducing *B. fragilis*, it was adequate to correct the abnormalities (Troy and Kasper 2010). Additionally, a *B. fragilis* strain isolated from a baby's feces was shown to have the capability to improve phagocytosis, improve IL-10 production, and increase CD80 and CD86 of M1 macrophage cell surface expression (Deng et al. 2016; Landuyt et al. 2021). There are proposals of developing *B. fragilis* as a probiotic because of its beneficial effect on its host (Deng et al. 2016). Other beneficial effects of other *Bacteroides* species are from *B. dorei*, *B. xylanisolvens*, *B. acidafaciens*, *B. ovatus*, and *B. uniformis*. These *Bacteroides* species have been proposed as live biotherapeutic products (LBP) or are also called next-generation probiotics (NGP). Generally, LBP/NGP is still under study and has no history yet of being used as a probiotic, unlike traditional probiotics (*Lactobacillus* and *Bifidobacterium*) with long and successful industrial use. Moreover, traditional probiotics are commonly used as food and supplements, but LBP/NGP is still being studied for a possible preclinical mode of action, pharmacokinetics, and pharmacodynamics (O'Toole et al. 2017).

Another example is *B. xylanisolvens* which was reported to possess immunomodulatory properties and produce short fatty acids (propionate and succinate) in the gut (Ulsemer et al. 2012). The research has shown that pasteurized *B. xylanisolvens*

increases the level of TNF $\alpha$ -specific IgM antibodies and promotes a robust response to cancer treatment.

Another beneficial relationship between mammalian host and *Bacteroides* that is well documented is *B. acidafaciens* which increases the production of IgA in the large intestine by increasing IgA<sup>+</sup> B cells (Yanagibashi et al. 2013). *B. uniformis*, on the other hand, when administered to obese mice, was shown to have beneficial effects in preventing obesity by significantly modifying total body weight. Further experiments have shown that the continuous administration of *B. uniformis* substantially reduces fat micelles in enterocytes and reduces leptin levels, serum cholesterol and triglyceride levels, which indicates that *B. uniformis* has a positive effect on the hepatic steatosis. Lastly, mice fed with the recombinant strain of *B. ovatus* that produces TGF- $\beta$ 1 can prevent or treat acute colitis (Gauffin Cano et al. 2012).

In another study, *B. fragilis* was shown to produce a capsular polysaccharide that helps protect a person from nerve demyelination (Ochoa-Reparaz et al. 2010). Furthermore, pediatricians analyzed the guts of 405 infants and found that those infants with *Bacteroides*-dominant microbiota were positively correlated with higher cognitive and language performance (Tamana et al. 2021). *B. fragilis* was even shown to produce large quantities of  $\gamma$ -aminobutyric acid (GABA), a neurotransmitter that influences behavior and response to stress (Strandwitz et al. 2019).

### *Bacteroides as a disease-causing organism*

Although *Bacteroides* is known to be a mutualist or commensal when in the gut, it is equally known to be pathogenic when it escapes the gut due to a breached or compromised mucosal wall of the intestine. This implies that location is pivotal. It has caused abscesses in the liver, abdominal cavity, brain, pelvis, lungs, perirectal area, gangrenous appendicitis, diverticulitis, inflammatory bowel disease, and bacteremia. There was even a study with 365,490 healthcare-related infections, which revealed that *Bacteroides* spp. is on the top 15 most common pathogen that causes nosocomial infections (Wexler 2007; Brook 2016; Weiner et al. 2016). Among the *Bacteroides* species, *B. fragilis* is identified to be the most virulent, with *B. ovatus*, *B. thetaiotamicron*, and *B. vulgatus* to be clinically important species (Majewksa et al. 2021). It is interesting because *B. fragilis* and *B. thetaiotamicron* are considered beneficial and at the same time, pathogenic (Mazmanian and Kasper 2006). The beneficial effects of *B. fragilis* are strain dependent. There are other strains of *B. fragilis* that produce fragilysin toxin associated with colorectal cancer (Cheng et al. 2020). Additionally, *B. fragilis* has been isolated from 63%–80% of *Bacteroides*-related infections. Conversely, *B. thetaiotamicron* accounts for 13%–17% of *Bacteroides* infection cases (Cerdeño-Tárraga et al. 2005). Moreover, some strains of *B. fragilis* and *B. thetaiotamicron* were found to have genes that encode C10 proteases. These *Bacteroides* C10 proteases were identified to be homologous to the C10 cysteine protease of *Streptococcus pyogenes* (Thornton et al. 2012). A genetic exchange may have caused such homology via HGT between the two bacteria. There are recent documentation of other *Bacteroides* species associated with infection such as *B. dorei* and *B. pyogenes*. An invasive strain of *B. dorei* was isolated from a patient and shown to have contributed to the patient's mycotic aortic aneurysm. This report suggests caution in using *B. dorei* for biotherapeutic purposes, as some have already proposed (O'Toole et al. 2017; Matsuoka et al. 2021). This is very interesting because *B. dorei* was considered to play a probiotic role in the gut (Zafar and Saier 2018). *B. pyogenes* can cause bacteremia secondary to cat and dog bites, prosthetic joint, and amputation infections (Majewksa et al. 2021). Moreover, *B. vulgatus* and *B. ovatus*

were associated with possible increased incidence of Crohn's and Coeliac diseases (Zafar and Saier 2018).

A very important characteristic of *Bacteroides* is their extensive ability for HGT. A study has shown that *Bacteroides* species engage more frequently in HGT than other bacterial species in the gut. *Bacteroides* displayed an average HGT frequency range of 0.85%–2.3% for 10-kb+ HGT and 6.0%–10.1% for 500-bp+ HGT. Additionally, two other gut bacteria: *Bifidobacterium longum* and *Akkermansia muciniphila* have a HGT frequency of 0.04% and 0.06% for 10-kb+ HGT and 0.81% and 1.64% for 500-bp+ HGT, respectively (Groussin et al. 2021). *Bacteroides* HGT is so extensive that it was reported to have performed interspecies and even interfamily HGTs – the most likely mechanism that *Bacteroides* gets other bacteria to acquire antibiotic resistance (Coyne et al. 2014). In the gut, *Bacteroides* serves as an antibiotic resistance gene reservoir that transfers those resistance genes to transient bacteria – those that just pass through or do not colonize the gut (Huddlestone 2014).

Quick genetic exchanges happen by conjugation as the favored mechanism of HGT due to the protection it provides to the DNA during transfer (Arber 2014). In another study, *B. fragilis* was shown to have a novel HGT mechanism that allows them to immediately insert very large segments of *Bacteroides* chromosomal DNA. Since large segments are transferred instead of multiple small segments, it allows *Bacteroides* to rapidly adapt to changing host GIT environment (Husain et al. 2017). It was also observed that most genes that contribute to strain diversity are in regions more likely to be acquired by HGT and spread out in multiple elements such as from phage, conjugative plasmids, and conjugative transposons (Coyne et al. 2014).

### Fecal water pollution

Bodies of water are one of the primary components of our ecosystem that play a fundamental role in the existence of all biological systems. For humans, water plays an important role in the fields of agriculture, livestock, forestry, industry, fisheries, aquaculture, energy production, recreation, and others (Effendi 2016; Ogunkunle et al. 2016). But for the simple yet vital part of it, water is an essential component of cellular homeostasis of the human body (Popkin et al. 2010). Unfortunately, there are reports of massive water pollution/contamination globally (Yan et al. 2015; Maceda-Veiga et al. 2017; Noorhosseini et al. 2017; Ruzol et al. 2017). Water pollution is defined as the direct/indirect and sudden or gradual negative alteration of water's physico-chemical and biological properties, which deteriorates water quality. Consequently, it becomes a hazard to the health of any organism and makes it unfit to its other assigned uses, incurring economic loss and social instability. Pollutants can be natural contaminants, such as those due to weathering and soil leaching which are gradual processes rather than anthropogenic sources that are rapid and usually on the rise and associated with industrial, agricultural, and domestic effluents (Ogunkunle et al. 2016). There are two types of water pollutant that affect water quality. These are the non-point sources (NPS) and point sources (PS). In NPS, pollutants enter the water system from unidentified diffuse sources and are difficult to control, like stormwater runoff and domestic and wild animal defecation. Alternatively, PS are impurities that enter the water system from an easily identifiable location via a direct route like effluents from sewage treatment plants, industrial effluents, and municipal wastes (Ahmed et al. 2005; Jamwal et al. 2011). An increasingly concerning water contaminant that threatens freshwater systems is fecal pollution, as it may cause the spread of microbial pathogens and increase the nutrient load of water (Fan et al. 2017). Contaminated feces from infected animal sources may contain various pathogenic microorganisms like *Escherichia coli*, *Campylobacter*,

*Salmonella*, *Shigella*, *Yersinia*, *Pasteurella*, *Francisella*, *Vibrio*, *Entamoeba histolytica*, *Blastocystis*, *Giardia*, and various viruses, such as adenoviruses, enteroviruses, and rotaviruses. Their release in freshwater bodies leads to waterborne diseases (Savichtcheva and Okabe 2006; Woodall 2009; Hampson et al. 2010; Efstratiou et al. 2017).

A major source of fecal pollution is from anthropogenic activities. It comes from agricultural, industrial, municipal, and household wastes (Li et al. 2021a). Poor fecal waste management in farms, inefficient discharge of water from treatment plants (Diaz-Gavidia et al. 2022), and reuse of municipal and domestic wastewater in agricultural irrigations (Al Hamedi et al. 2023) are the causes of widespread fecal pollution. However, efforts have been made to solve this challenge. Yet, the situation of fecal pollution lingers because of our inability to pinpoint the sources of the pollutant – a key step in mitigating the fecal pollution problem (Xue et al. 2018). If the point of origin of fecal contamination and pathogens is precisely and accurately known whether human or animal, then the proper government policy, management, minimization of public health risks associated with fecal contamination, and remediation efforts can be apportioned more efficiently and effectively (Ahmed et al. 2005; Savichtcheva and Okabe 2006). Thus, a relatively new technique that answers the question of reliably pinpointing the source of the fecal contamination by using fecal bacteria and determining how much of each source contributes to the contamination is called microbial source tracking (MST) (Harwood et al. 2014).

### Microbial source tracking

MST, the same as fecal source tracking and bacterial source tracking, is based on fecal microorganisms being strongly associated with their host species. Moreover, it supports the idea that some identified attributes or patterns (i.e., antibiotic resistance patterns, 16S rRNA gene sequences) of the host-associated microorganisms can be used as markers for tracing the fecal contamination back to its host (Harwood et al. 2014). It is a relatively new approach to discriminate, pinpoint, and quantify human and other fecal contamination that may exist in the environment and is usually applied in aquatic environments (Odagiri et al. 2015; Mattioli et al. 2017). In short, MST is about identifying the specific animal source of a given fecal contaminant. This contrasts with the traditional culture-based method that is limited to determining the presence of a fecal contaminant (Rivera and Rock 2011).

Globally, different MST methods have been used to accomplish the goal of MST, which is to determine fecal pollution sources so that the needed socio-political and environmental actions can be undertaken. Some MST research was conducted in Virginia, USA, in which antibiotic resistance patterns and carbon utilization profiles were used to find out whether there was fecal pollution in their waters and to trace its specific sources (Wiggins 1996; Hagedorn et al. 2003). The biochemical fingerprinting MST method was used in Queensland, Australia, to identify the sources of *E. coli* and *Enterococci* in Eudio, Creek (Ahmed et al. 2005). The human specific *HF183 Bacteroides* MST marker was used to assess the extent of human fecal pollution in two freshwater canals in Ghent, Belgium. Conversely, a similar assessment but on pig fecal pollution was done in Brittany, France (Mieszkina et al. 2009). Fatty acid methyl ester (FAME) profiling analysis was used in Pasadena, USA, to source-track the fecal coliforms (Duran et al. 2006). In Michigan, USA, MST method using the  $\beta$ -glucuronidase gene was used to know whether household pets have a significant input in the fecal pollution in the surrounding surface waters (Ram et al. 2007). In France, two library-independent MST methods were used (Bacteroidales 16S rRNA gene marker

detection and F-specific RNA bacteriophage genotyping) to source track the fecal pollution in a French estuary (Gourmelon et al. 2007). MST was also used in Florida, USA, to evaluate the effect of stormwater concerning water quality using BOX-PCR (Brownell et al. 2007). In Nicoya, Costa Rica, MST was used to determine the water quality of shellfish harvesting waters using both viral and bacterial based MST markers (Symonds et al. 2017). Other MST research was conducted in Korea, using F+ RNA coliphages to source track fecal pollution in Metropolitan Seoul and Gyeonggi province (Lee et al. 2011), Lake Taihu in China (Hagedorn and Xinqiang 2011; Vadde et al. 2019), and Dargle River and its tributaries in Dublin, Ireland (Ballesté et al. 2020). Interestingly, MST was even used to test for fecal contamination in fresh produce like tomatoes, jalapeño peppers, and cantaloupe and to even determine the sources of the fecal

contaminant (Lee and Lee 2012; Ravaliya et al. 2014). This showed that MST could also be used in other aspects of research outside its usual or traditional context, which is for assessing environmental water fecal pollution. These studies conducted globally show the scale at which MST has been successfully used to monitor fecal pollution, particularly in developed nations. Unfortunately, its application in many developing countries is still limited (Sommark et al. 2018a).

MST methods are either library-dependent (LDM) or library-independent (LIM) and are analyzed either based on their phenotypic and/or genotypic characteristics, as shown in Table 2.

**Table 2: Techniques under library-dependent and library-independent MST methods**

Library-dependent		Library-independent	
Culture-based		Culture-independent	
Phenotypic (Biochemical)	Genotypic	Phenotypic or Genotypic	Molecular
<ul style="list-style-type: none"> <li>Antibiotic resistance (ARA and MARA)</li> <li>Carbon utilization</li> <li>FAME profiling</li> </ul>	<ul style="list-style-type: none"> <li>Rep-PCR</li> <li>RAPD</li> <li>AFLP</li> <li>PFGE</li> <li>Ribotyping</li> <li>DGGE</li> <li>LH-PCR</li> <li>T-RFLP</li> </ul>	<ul style="list-style-type: none"> <li>Bacteriophage</li> </ul>	<ul style="list-style-type: none"> <li>Host-specific viral PCR</li> <li>Host-specific bacterial PCR</li> </ul>

LDM entails creating a comprehensive database of isolates from known sources that will serve as the basis for cross-referencing. Firstly, fecal samples are collected from potential or suspected sources (i.e., humans, cattle, swine, etc.) of water contamination. Then, bacteria from feces are isolated and subjected to either phenotypic or genotypic analysis to create a pattern referred to as a fingerprint used for cross-referencing. Phenotypic analysis may include antibiotic resistance analysis (ARA), multiple antibiotic resistance analysis (MARA), carbon source utilization analysis, and FAME profiling. Conversely, genotypic analysis includes patterns in repetitive DNA sequences (Rep-PCR), random amplification of polymorphic DNA (RAPD), amplified fragment-length polymorphism (AFLP), pulse-field gel electrophoresis (PFGE), ribotyping, denaturing-gradient gel electrophoresis (DGGE), length heterogeneity PCR (LH-PCR), and terminal restriction fragment-length polymorphism analysis (T-RFLP) (Hagedorn et al. 2003; Meays et al. 2004; Duran et al. 2006; Ballesté and Blanch 2010). However, a major limitation in LDM is that it is highly tedious to make a reliable database because it requires numerous isolates to be tested and even then, the minimum number of isolates needed to construct a reliable library has not been fully established (Ahmed et al. 2005).

Alternatively, LIM does not require the construction of a database but depends on the concept that bacteria or viruses taken directly from the environment come from species-specific host/s or sources of fecal contamination; thus, a library is not needed for cross-referencing. A significant advantage of LIM, as opposed to LDM, is that it saves time and resources as it removes the need to construct a library starting with the isolation, purification, and culture of microorganisms. In LIM, detection of MST-specific markers (or simply MST markers) is used. There have been various MST markers developed to discriminate human fecal from other fecal animal sources like viral MST markers (Wong et al. 2012), bacterial MST markers, and recently, the use of mitochondrial DNA (mtDNA) as MST markers (He et al. 2016). The most common type of MST marker is the use of *Bacteroides* spp. as indicators of the type of host animal serving as the source of fecal pollution. *Bacteroides* MST

markers are designed to target specific diagnostic sequences within the *Bacteroides* 16S rRNA gene present in feces from different animals (Layton et al. 2006).

#### *Bacteroides*-based MST markers

*Bacteroides*-based MST marker is the most extensively used MST marker. Due to the anaerobic nature of *Bacteroides*, it has minimal capacity for growth outside the GIT of its host. This makes it ideal for MST because it provides a more accurate count in terms of the degree of fecal pollution. It has shown a high level of host specificity as *Bacteroides* adapt to the GIT conditions of its host and has even co-evolved with its host species (Bernhard and Field 2000). Also, the traditional FIB only represents less than 2% of the gut microbiota. FIB can remain viable and multiply in environmental waters, giving a false alarm of fecal pollution (Zheng and Shen 2018). Lastly, there is a strong positive correlation that the presence of *Bacteroides* also indicates the presence of fecal pathogens like *E. coli*, *Salmonella*, and *Campylobacter*, as shown in Table 3, as opposed to detecting the presence of traditional FIB that does not show a similar correlation. Therefore, due to all those characteristics, many LIM are designed to target specific sequences within the *Bacteroides* 16S rRNA gene to discriminate the different sources of contamination by PCR. Some of the known *Bacteroides*-based MST markers are shown in Table 4.

**Table 3: Studies reporting strong positive correlation between the presence of *Bacteroides* and the presence of fecal pathogens**

Analysis	Studies
<i>Bacteroides</i> and fecal coliform	Schriewer et al. 2010; Villemur et al. 2015
<i>Bacteroides</i> and <i>Escherichia coli</i>	Ahmed et al. 2008; Mauffret et al. 2012; Nshimiyimana et al. 2014
<i>Bacteroides</i> and enteroviruses	Noble et al. 2006; Sauer et al. 2011
<i>Bacteroides</i> and <i>Enterococcus</i>	Eichmiller et al. 2013; Gordon et al. 2013
<i>Bacteroides</i> and <i>Salmonella</i> , <i>Campylobacter</i>	Schriewer et al. 2010; Stea et al. 2015

<i>Bacteroides</i> and <i>Giardia</i> , <i>Cryptosporidium</i>	Staley et al. 2012; Marti et al. 2013
--	---------------------------------------

**Table 4: Known *Bacteroides*-based MST markers and its target host**

Specific target host	MST marker/s
Human	<i>HF183</i> (Bernhard and Field 2000)
	<i>HuBac</i> (Layton et al. 2006)
	<i>BacHum</i> (Kildare et al. 2007)
	<i>BacH</i> (Reischer et al. 2007)
	<i>HumM2</i> (Shanks et al. 2009)
	<i>HumM3</i> (Shanks et al. 2009)
Pig	<i>PigBac-1</i> and <i>PigBac-2</i> (Okabe et al. 2007)
	<i>Pig-1-Bac</i> and <i>Pig-2-Bac</i> (Mieszkin et al. 2009)
Cow	<i>BacCow</i> (Kildare et al. 2007)
	<i>BoBac</i> (Layton et al. 2006)
Dog	<i>BacCan</i> (Kildare et al. 2007)
Horse	<i>HoF597</i> (Dick et al. 2005)
Muskrat	<i>MuBa01</i> (Marti et al. 2011)
General Ruminant	<i>Rum-2-Bac</i> (Mieszkin et al. 2010)
	<i>BacR</i> (Reischer et al. 2006)
Chicken	<i>ChickenBac</i> (Kobayashi et al. 2013)
Duck	<i>DuckBac</i> (Kobayashi et al. 2013)
Canada Goose	<i>CGOF1-Bac</i> (Fremaux et al. 2010)
	<i>CGOF2-Bac</i> (Fremaux et al. 2010)
General Feces	<i>AllBac</i> (Layton et al. 2006)

The effectivity of MST and its *Bacteroides*-based markers have been tried and tested in several research globally since their development. *HF183* gene marker was used as a tracer in discriminating between human and ruminant sources of fecal pollution in New River, Pasco County, Florida, USA (Chase et al. 2012), in Lake Pontchartraine, Louisiana, USA (Xue et al. 2018), in Nijo River, Kenya (Jenkins et al. 2009), in Danube River (Kirschner et al. 2017) and Daoulas River catchment, Brittany, France (Mauffret et al. 2012). It was also used in determining whether sand and other submerged sediments in the waters of Lake Superior, Minnesota, USA can also serve as a reservoir of fecal pollution and thus can be an alternative source for MST sampling other than water samples (Eichmiller et al. 2013). Conversely, *HF183* was the MST marker used to establish the positive correlation of MST markers with enterococci in the Gulf of Mexico waters (Gordon et al. 2013). *HuBac* marker was used in the coastal water of Puerto Rico and Trinidad and Tobago to assess fecal pollution (Bachoon et al. 2010). In Ireland, the human MST marker *BacHum* was used to assess the water quality of Mattock River catchment (Flynn et al. 2016), Ware and Oyster Creek, North Carolina, USA (Gonzalez et al. 2012), Sava River Basin that borders three countries such as Serbia, Bosnia and Herzegovina, and Croatia (Vrzel et al. 2016), and Monterey Bay, central California coast, USA (Schriewer et al. 2010). *BacH* was developed in 2006 to detect and discriminate human fecal pollution in the Northern Alps, Austria (Reischer et al. 2006). In Israel, *BacH* marker was the human MST marker used to assess the water quality of Karst springs found in the Sea of Galilee (Ohad et al. 2015), alpine springs in Austria (Reischer et al. 2007), coast of Odisha, Bay of Bengal (Odagiri et al. 2015) and in Cornwallis watershed, Nova Scotia, Canada (Ridley et al. 2014).

*Pig-2-Bac* is often used to detect and source track pig fecal pollution (Bae and Wuerz 2015). *Pig-2-Bac* was the pig marker used to assess the freshwater quality of Taige and Taihu Lake in China (He et al. 2016), in Danube River (Kirschner et al., 2017), Daoulas catchment, France (Mieszkin et al. 2009; Mauffret et al. 2012), in Sava River Basin in Central Europe (Vrzel et al. 2016), and Tha Chin watershed in Thailand (Somnark et al. 2018b). MST marker assessment showed that *BacCow*, as the cow MST marker, was used for India (Odagiri et al. 2015) and Thailand

(Somnark et al. 2018b). It was also used to identify fecal sources of contamination in the Njoro River watershed, Kenya (Jenkins et al. 2009), surface waters of the central California coast, USA (Schriewer et al. 2010), and coastal waters in Odisha, India (Odagiri et al. 2015). Alternatively, *BoBac* cow MST marker was used to assess the NPS of fecal pollution in stream waters of Brandywine Creek that pass by the states of Pennsylvania, and Delaware, USA (Duris et al. 2011), Harris Neck estuarine marshes in Georgia, USA (Markand et al. 2011), pond waters, the primary source of water for hygiene purposes in Bangladesh (Knappett et al. 2011), and sediment water in Saginaw Bay, Michigan, USA (Oun et al. 2017). Cow MST marker *BacBov* was used to assess the water quality in the Grand River watershed, southwestern Ontario, Canada, and in Mattock River catchment that supplies water to 7% arable land in Ireland (Flynn et al. 2016). *Rum2Bac* is a specific marker for detecting fecal contaminants of generally all ruminants that includes domesticated animals such as cows, carabaos, sheep, and goats and nondomesticated ones like giraffes and elks. Ruminants are mammals composed of four gastric compartments namely: rumen, reticulum, omasum, and abomasum. The first three mentioned stomachs are collectively known as pro-ventriculus as they share a common function. They contain bacteria that break down ingested cellulose from grasses that ruminants usually eat (Teixeira et al. 2009). Thus, *Rum2Bac* is commonly used to screen for the presence of feces from other ruminants other than that of cows. *Rum2Bac* has been used in France in their development of MST tools to identify the origins of fecal pollution in bathing and shellfish harvesting waters (Gourmelon et al. 2010; Mauffret et al. 2013) and in assessing the water quality in Daoulas catchment, Justicou, Pen an Traon, and La Fresnaye, Brittany, France (Mauffret et al. 2012; Jardé et al. 2018). *Rum2Bac* (with other markers) was used in a study conducted in South Fork Broad River, Georgia, USA, to show the relationship between MST markers and FIB. They could establish a positive correlation between Shiga-toxin, *Campylobacter*, and *Rum2Bac* MST marker (Bradshaw et al. 2016).

*BacCan* is the only known/developed *Bacteroides*-based MST marker for dogs. It is the MST marker used to determine the water quality and source track the waters of Ningi Creek, Queensland, Australia (Ahmed et al. 2008), coastal surface waters, central California, USA (Schriewer et al. 2010), different dog fountains and ponds across Calgary, Alberta, Canada (Tambalo et al. 2012); coast of Odisha, part of Bay of Bengal, India (Odagiri et al., 2015), and Gulf of Nicoya, Costa Rica (Symonds et al. 2017). In 2011, an MST marker for muskrat (*Ondatra zibethicus*) was developed to determine the presence and extent of the muskrat's fecal pollution contributions to South Nations River, Canada. *MuBa01*, when tested, showed a sensitivity of 66% and specificity of 100% (Marti et al. 2011). Muskrats are wide-ranging semi-aquatic mammals common to North America and considered a normal and very common feature of the wetlands. They have very high fecundity, producing 2–3 offspring per year. Additionally, muskrats born in that year can also breed within that year due to their fast growth (Sadowski and Bowman 2021).

*AllBac* is primarily a screening or preliminary MST marker to first check the presence (if any) of fecal contamination in a body of water. It is the initial test before performing or using more host-specific MST markers. Such as in the research by Stapleton et al. (2009), in which they used *AllBac* to first detect fecal contamination in the recreational bathing waters in the UK. A similar study was conducted in France, but other than bathing waters, it also included shellfish harvesting waters and the Daoulas catchment (Gourmelon et al. 2010; Mauffret et al. 2012). The same was done in rural and urban watersheds in

Canada by using *AllBac* to monitor fecal pollution (Ridley et al. 2014; Stea et al. 2015), in the Danube River (Mayer et al. 2016; Kirschner et al. 2017), in the Sava River Basin, Central Europe (Vrzal et al. 2016), and in the beach area of Duluth-Superior Harbor, Minnesota, USA (Zhang et al. 2016). In Bangladesh, *AllBac* was used to generalize that there are other sources of fecal contamination other than humans and cows in several pond waters used for hygiene purposes (Knappett et al. 2011).

As can be seen, most of the markers are for mammalian fecal samples. However, *Bacteroides* has been identified in other warm-blooded and cold-blooded vertebrates such as ducks, geese, frogs, and toads. Unfortunately, for avian MST, only three MST markers have been developed. Many migratory waterfowls such as ducks and geese and non-migratory birds like chickens are also carriers of pathogens. Hence, the three avian-specific MST markers developed are insufficient in better monitoring and managing water pollution. After development, *ChickenBac* and *DuckBac* were immediately used to check the water quality by MST in Kamogamo River, Toyohira River, and Oono pond, Hokkaido, Japan, and showed an abundance of fecal contamination from ducks (Kobayashi et al. 2013). Alternatively, Canada's geese population was estimated to be at 3 million and often occupy regions within urban settings. For that reason, the *CGOF* MST markers were developed to determine the extent of fecal pollution that Canadian geese may have contributed. True enough, using the developed *CGOF* markers, water samples from Wascana Lake detected high prevalence and abundance of *Bacteroides* markers (Fremaux et al. 2010).

However, there have been reports of a cross-reaction of *Bacteroides*-based MST gene markers with other fecal samples. There are numerous cases where several MST markers supposed to be for a specific host would vary in their specificity. The two MST markers for pigs (*Pig-1-Bac* and *Pig-2-Bac*) would usually vary in their specificity depending on the geography (Mieszkin et al. 2009; Heaney et al. 2015). Another would be among the different human associated *Bacteroides* MST markers. Although *HF183* is the most widely used, it is not always the best performing MST marker. *HF183* was identified as a high-performing marker in California, USA, and Australia. However, the MST marker *BachHum* was the better performing MST marker in Kenya and India. The 16S rRNA gene is a highly conserved region among bacteria. Thus, it is for this reason that cross-reactivity happens when this gene is amplified in MST. Therefore, it is important that marker sensitivity and specificity are established before performing actual MST research as they are the two major performance indicators of an MST marker and determine which MST marker to use (Ahmed et al. 2016).

MST sensitivity is defined as the percentage of samples from a targeted host positive for the chosen marker. Likewise, the greater the sensitivity of a marker, the closer it is to 100% host sensitivity. It is empirically measured by testing a collection of fecal samples from the target host, or it is simply expressed in Equation 1, mathematically (Nshimiyimana et al. 2014). Currently, there is no consensus on the number of samples, type of sample, and sample volume needed to determine the sensitivity of a marker. Unfortunately, there is also no universal benchmark for sensitivity, but a value of > 80% is acceptable for MST use (Ahmed et al. 2016). Also, sensitivity varies due to geographical variations. This means that an MST marker with high sensitivity used in a given location does not necessarily mean that it will also have an increased sensitivity if used in a different location, because it has precedence from a previous site as being a highly sensitive marker. Thus, MST marker sensitivity should always be verified in every geographic region before performing a study (Harwood et al. 2014).

$$\text{Equation 1: Sensitivity} = \left( \frac{\text{TP}}{\text{TP} + \text{FN}} \right) \times 100$$

Where: TP = true positive  
FN = false negative

Conversely, specificity is the number of non-target host samples that test negative for the assayed marker and mathematically shown in Equation 2 (Nshimiyimana et al. 2014). Much research worldwide shows that most MST gene markers have variable specificity in different geographic areas. Studies have shown that an MST marker specific to a particular organism would cross-react to different organism/s. Unfortunately, like sensitivity, there is no recognized standard about the number of samples for specificity testing. However, statistically, analyzing as many non-target fecal samples as possible will increase confidence (Harwood et al. 2014).

$$\text{Equation 2: Specificity} = \left( \frac{\text{TN}}{\text{TN} + \text{FP}} \right) \times 100$$

Where: TN = true negative  
FP = false positive

Ultimately, sensitivity and specificity vary with geographic area and time, highlighting the need for local validation. Therefore, before conducting any MST study, it is critical to first establish sensitivity and specificity, and validate MST marker accuracy using Equation 3 (Nshimiyimana et al. 2014). Moreover, since the number of samples to reliably establish sensitivity and specificity is currently not yet standardized, some recent studies have reported that between 12–20 samples is acceptable (Reischer et al. 2006; Kildare et al. 2007; Haugland et al. 2010; Ebentier et al. 2013).

$$\text{Equation 3: Accuracy} = \left( \frac{\text{TP} + \text{TN}}{\text{TP} + \text{FP} + \text{TN} + \text{FN}} \right) \times 100$$

Where: TN = true negative  
TP = true positive  
FP = false positive  
FN = false negative

#### Future research avenues

*Bacteroides*, either as a “good” or “bad” bacterium, do not receive much attention as opposed to *Lactobacillus* and *Bifidobacterium* for being “good bacteria” and different fecal coliform bacteria as being “bad bacteria.” At times, they are



described simply as gut microorganisms. In Europe, the European-wide surveillance on antibiotic resistance of *Bacteroides* is conducted annually, to determine the current state of antimicrobial resistance (AMR) among the *Bacteroides* group (Kierzkowska et al. 2020; SÓki et al. 2020). Thus, more research on the AMR of *Bacteroides* must also be conducted in other countries in a similar way that other bacteria such as the AMR of *E. coli*, *Staphylococcus*, and *Klebsiella* are being extensively monitored. Another avenue for *Bacteroides* research is on *Bacteroides* from animals. Most of the research on *Bacteroides* is human-centered. However, more than half of the world's antibiotic usage is for animal husbandry, but ironically, a meager 10% of publications on antibiotic resistance are about animals, and the rest are on humans (He et al. 2020).

There are many animals globally that contribute to fecal water pollution, and yet there are only a handful of MST markers developed. Understandably, the only MST markers made were for those that contribute large volumes of feces into the aquatic environment, such as humans and domesticated farmed animals. However, to create a long-term water conservation policy, more MST markers should be developed because other non-domesticated animals are also carriers of virulent microorganisms. Also, even among the domesticated animals, there are still many of which are without a developed MST marker like goats, sheep, and poultry like turkey.

Different kinds of migratory flying animals like pigeons, sea gulls, pelicans, herons, geese, swans, and bats have been associated as disease carriers of *Borrelia* (Lyme disease), West Nile virus (viral encephalitis), *Mycobacterium avium/genevense* (mycobacteriosis), influenza A virus (avian flu), Nipah virus, rabies virus, and SARS coronavirus. These migratory flying animals come by hundreds to thousands bringing with them pathogens. However, there are only two *Bacteroides*-based MST markers developed for migratory flying animals, which include for ducks (*DuckBac*) and for Canada geese (*CGOF*) (Reed et al. 2003; Calisher et al. 2006). Therefore, more MST markers should be developed for migratory flying animals.

Another aspect that can be enhanced is the accuracy of the MST markers developed. Note that diet serves as a strong evolutionary pressure on the gut microbiome. For example, *DuckBac* marker was developed using wild migratory ducks; hence, it may or may not be sensitive or specific enough if it is used for farmed ducks due to obvious differences in the diet of wild and farmed ducks. Thus, this also brings the issue of marker cross-reactivity. Therefore, it may be time to use a different approach in developing *Bacteroides*-based markers to increase specificity and sensitivity, like using other genes (housekeeping genes, virulence genes, antibiotic resistance genes).

In conclusion, *Bacteroides* are an interesting group of bacteria found in the gut of many animals occupying an important ecological niche of the gut microbiome. Most of their attributions are on their role in the gut microbiome. However, their use on MST has now become a critical tool in helping to curb the problem of fecal water pollution. Lastly, it is noticeable that most of the countries that perform MST are highly developed countries like Canada, the USA, China, Japan, and France. After emphasizing the advantage of MST over the traditional FIBs, it is time for other countries, through their scientists, to explore the use of MST in monitoring water pollution. There is an imminent need for collaboration and cross-disciplinary research between countries' environmental managers, microbial ecologists, and government environmental policymakers to transfer and share the MST technology so that its use can be a norm.

## ACKNOWLEDGMENTS

This study was financially supported by the National Research Council of the Philippines (Project No. E-249) and the Philippine Council for Industry, Energy and Emerging Technology Research and Development.

## CONFLICT OF INTEREST

The authors declare that there is no conflict of interest.

## REFERENCES

- Ahmed W, Hughes B, and Harwood VJ. Current status of marker genes of *Bacteroides* and related taxa for identifying sewage pollution in environmental waters. *Water (Switzerland)* 2016; 8(6). <https://doi.org/10.3390/w8060231>
- Ahmed W, Neller R, and Katouli M. Host species-specific metabolic fingerprint database for tracking the sources of faecal contamination in surface waters. *Applied and Environmental Microbiology* 2005; 71: 4461–4468. <https://doi.org/10.1128/AEM.71.8.4461-4468.2005>
- Ahmed W, Powell D, Goonetilleke A, and Gardner T. Detection and source identification of faecal pollution in non-sewered catchment by means of host-specific molecular markers. *Water Science and Technology* 2008; 58(3): 579–586. <https://doi.org/10.2166/wst.2008.436>
- Al Hamed FH, Kandhan K, Liu Y, Ren M, Jaleel A, and Alyafei MAM. Wastewater irrigation: A promising way for future sustainable agriculture and food security in the United Arab Emirates. *Water (Switzerland)* 2023; 15(12), 1–18. <https://doi.org/10.3390/w15122284>
- Arber W. Horizontal gene transfer among bacteria and its role in biological evolution. *Life* 2014; 4(2), 217–224. <https://doi.org/10.3390/life4020217>
- Bachoon DS, Markand S, Otero E, Perry G, and Ramsubhag A. Corrigendum to “Assessment of non-point sources of fecal pollution in coastal waters of Puerto Rico and Trinidad” [*Marine Pollution Bulletin* 60 (2010) 1117–1121]. *Marine Pollution Bulletin* 2010; 60(12): 2358. <https://doi.org/10.1016/j.marpolbul.2010.11.012>
- Bäckhed F, Ley RE, Sonnenburg JL, Peterson DA, and Gordon JI. Host-bacterial mutualism in the human intestine. *Science* 2005; 307(5717): 1915–1920. <https://doi.org/10.1126/science.1104816>
- Bae S, and Wuertz, S. Decay of host-associated Bacteroidales cells and DNA in continuous-flow freshwater and seawater microcosms of identical experimental design and temperature as measured by PMA-qPCR and qPCR. *Water Research* 2015; 70: 205–213. <https://doi.org/10.1016/j.watres.2014.10.032>
- Bakir MA, Kitahara M, Sakamoto M, Matsumoto M, and Benno Y. *Bacteroides finegoldii* sp. nov., isolated from human faeces. *International Journal of Systematic and Evolutionary Microbiology* 2006a; 56(5): 931–935. <https://doi.org/10.1099/ijs.0.64084-0>
- Bakir MA, Kitahara M, Sakamoto M, Matsumoto M, and Benno Y. *Bacteroides intestinalis* sp. nov., isolated from human faeces. *International Journal of Systematic and Evolutionary*

- Microbiology 2006b; 56(1): 151–154. <https://doi.org/10.1099/ijs.0.63914-0>
- Ballesté E, and Blanch AR. Persistence of *Bacteroides* species populations in a river as measured by molecular and culture techniques. *Applied and Environmental Microbiology* 2010; 76(22): 7608–7616. <https://doi.org/10.1128/AEM.00883-10>
- Ballesté E, Demeter K, Masterson B, Timoneda N, Sala-Comorera L, and Meijer WG. Implementation and integration of microbial source tracking in a river watershed monitoring plan. *Science of the Total Environment* 2020; 736. <https://doi.org/10.1016/j.scitotenv.2020.139573>
- Benno Y, Watabe J, and Mitsuoka T. *Bacteroides pyrogenes* sp. nov., *Bacteroides suis* sp. nov., and *Bacteroides helcogenes* sp. nov., new species from abscesses and feces of pigs. *Systematic and Applied Microbiology* 1983; 4(3): 396–407. [https://doi.org/10.1016/S0723-2020\(83\)80024-1](https://doi.org/10.1016/S0723-2020(83)80024-1)
- Bernhard AE, and Field KG. A PCR assay to discriminate human and ruminant feces on the basis of host differences in *Bacteroides-Prevotella* genes encoding 16S rRNA. *Applied and Environmental Microbiology* 2000; 66(10): 4571–4574. <https://doi.org/10.1128/AEM.66.10.4571-4574.2000>
- Bradshaw J, Snyder BJ, Oladeinde A, Spidle D, Berrang ME, Meinersmann RJ, Oakley B, Sidle RC, Sullivan K, and Molina M. Characterizing relationships among fecal indicator bacteria, microbial source tracking markers, and associated waterborne pathogen occurrence in stream water and sediments in a mixed land use watershed. *Water Research* 2016; 101: 498–509. <https://doi.org/10.1016/j.watres.2016.05.014>
- Brook I. Spectrum and treatment of anaerobic infections. *Journal of Infection and Chemotherapy* 2016; 22(1): 1–13. <https://doi.org/10.1016/j.jiac.2015.10.010>
- Brownell MJ, Harwood VJ, Kurz RC, McQuaig SM, Lukasik J, and Scott TM. Confirmation of putative stormwater impact on water quality at a Florida beach by microbial source tracking methods and structure of indicator organism populations. *Water Research* 2007; 41: 3747–3757. <https://doi.org/10.1016/j.watres.2007.04.001>
- Calisher CH, Childs JE, Field HE, Holmes KV, and Schountz T. Bats: important reservoir hosts of emerging viruses. *Clinical Microbiology Reviews* 2006; 19(3): 531–545. <https://doi.org/10.1128/CMR.00017-06>
- Castellani A, and Chalmers AJ. *Manual of Tropical Medicine* 1919; 3rd ed. Williams, Wood and Co., New York
- Cerdeño-Tárraga AM, Patrick S, Crossman LC, Blakely G, Abratt V, Lennard N, Poxton I, Duerden B, Harris B, Quail MA, Barron A, Clark L, Corton C, Doggett J, Holden MTG, Larke N, Line A, Lord A, Norbertczak H, Ormond D, Price C, Rabbinowitsch E, Woodward J, Barrell B, and Parkhill J. Extensive DNA inversions in the *B. fragilis* genome control variable gene expression. *Science* 2005; 307(5714): 1463–1465. <https://doi.org/10.1126/science.1107008>
- Chase E, Hunting J, Staley C, and Harwood VJ. Microbial source tracking to identify human and ruminant sources of faecal pollution in an ephemeral Florida river. *Journal of Applied Microbiology* 2012; 113(6): 1396–1406. <https://doi.org/10.1111/jam.12007>
- Chassard C, Delmas E, Lawson PA, and Bernalier-Donadille A. *Bacteroides xylanisolvans* sp. nov., a xylan degrading bacterium isolated from human faeces. *International Journal of Systematic and Evolutionary Microbiology* 2008; 58(4): 1008–1013. <https://doi.org/10.1099/ijs.0.65504-0>
- Cheng WT, Kantilal HK, and Davamani F. The mechanism of *Bacteroides fragilis* toxin contributes to colon cancer formation. *Malaysian Journal of Medical Sciences* 2020; 27(4): 9–21. <https://doi.org/10.21315/mjms2020.27.4.2>
- Comstock LE. Importance of glycans to the host-*Bacteroides* mutualism in the mammalian intestine. *Cell Host and Microbe* 2009; 5(6): 522–526. <https://doi.org/10.1016/j.chom.2009.05.010>
- Coyne MJ, Zitomersky NL, McGuire AM, Earl AM, and Comstock LE. Evidence of extensive DNA transfer between *Bacteroidales* species within the human gut. *MBio* 2014; 5(3). <https://doi.org/10.1128/mBio.01305-14>
- Dela Rosa CJO, and Rivera WL. Identification of *Bacteroides* spp. from ducks using 16S rRNA gene PCR assay: prelude to its application in microbial source tracking. *Journal of Microbiology Biotechnology and Food Sciences* 2021; 1–7. <https://doi.org/10.15414/jmbfs.4101>
- Deng H, Li Z, Tan Y, Guo Z, Liu Y, Wang Y, Yuan Y, Yang R, Bi Y, Bai Y, and Zhi F. A novel strain of *Bacteroides fragilis* enhances phagocytosis and polarizes M1 macrophages. *Scientific Reports* 2016; 6(20): 1–11. <https://doi.org/10.1038/srep29401>
- Díaz-Gavidia C, Barría C, Weller DL, Salgado-Caxito M, Estrada EM, Araya A, Vera L, Smith W, Kim M, Moreno-Switt AI, Olivares-Pacheco J, and Adell AD. Humans and hoofed livestock are the main sources of fecal contamination of rivers used for crop irrigation: A Microbial source tracking approach. *Frontiers in Microbiology* 2022; 13. <https://doi.org/10.3389/fmicb.2022.768527>
- Dick LK, Bernhard AE, Brodeur TJ, Santo Domingo JW, Simpson JM, Walters SP, and Field KG. Host distributions of uncultivated fecal *Bacteroidales* bacteria reveal genetic markers for fecal source identification. *Applied and Environmental Microbiology* 2005; 71(6): 3184–3191. <https://doi.org/10.1128/AEM.71.6.3184-3191.2005>
- Duran M, Haznedarog BZ, and Zitomer DH. Microbial source tracking using host specific FAME profiles of fecal coliforms. *Water Research* 2006; 40: 67–74. <https://doi.org/10.1016/j.watres.2005.10.019>
- Duris JW, Reif AR, Olson LE, and Johnson HE. Pathogenic bacteria and microbial-source tracking markers in Brandywine Creek Basin, Pennsylvania and Delaware, 2009–10: U.S. Geological Survey Scientific Investigations Report 2011; 27pp.
- Ebentier DL, Hanley KT, Cao Y, Badgley BD, Boehm AB, Ervin JS, Goodwin KD, Gourmelon M, Griffith JF, Holden PA, Kelty CA, Lozach S, McGee C, Peed LA, Raith M, Ryu H, Sadowsky MJ, Scott EA, Domingo JS, Schiewer A, Sinigalliano CD, Shanks OC, Van De Werhorst LC, Wang D, Wuertz S, and Jay JA. Evaluation of the repeatability and reproducibility of a suite of qPCR-based microbial source tracking methods. *Water Research* 2013; 47(18): 6839–6848. <https://doi.org/10.1016/j.watres.2013.01.060>
- Eckburg PB, Bik EM, Bernstein CN, Purdom E, Dethlefsen L, Sargent M, Gill SR, Nelson KE, and Relman DA. Diversity of

- the human intestinal microbial flora. *Science* 2005; 308(5728): 1635–1638.
- Effendi H. River water quality preliminary rapid assessment using pollution index. *Procedia Environmental Sciences* 2016; 33: 562–567. <https://doi.org/10.1016/j.proenv.2016.03.108>
- Efstratiou A, Ongerth JE, and Karanis P. Waterborne transmission of protozoan parasites: Review of worldwide outbreaks - An update 2011–2016. *Water Research* 2017; 114: 14–22. <https://doi.org/10.1016/j.watres.2017.01.036>
- Eggerth AH, and Gagnon BH. The *Bacteroides* of human feces. *Journal of Bacteriology* 1932; 25(4). <https://doi.org/10.1128/jb.25.4.389-413.1933>
- Eichmiller JJ, Hicks RE, and Sadowsky MJ. Distribution of genetic markers of fecal pollution on a freshwater sandy shoreline in proximity to wastewater effluent. *Environmental Science and Technology* 2013; 47(7): 3395–3402. <https://doi.org/10.1021/es305116c>
- Fan L, Shuai J, Zeng R, Mo H, Wang S, Zhang X, and He Y. Validation and application of quantitative PCR assays using host specific Bacteroidales genetic markers for swine fecal pollution tracking. *Environmental Pollution* 2017; 231: 1569–1577. <https://doi.org/10.1016/j.envpol.2017.09.047>
- Ferrocino I, Di Cagno R, De Angelis M, Turrone S, Vannini L, Bancalari E, Rantsiou K, Cardinali G, Neviani E, and Coccolin L. Fecal microbiota in healthy subjects following omnivore, vegetarian and vegan diets: Culturable populations and rRNA DGGE profiling. *PLoS ONE* 2015; 10(6), 1–16. <https://doi.org/10.1371/journal.pone.0128669>
- Flynn RM, Deakin J, Archbold M, Cushnan H, Kilroy K, O’Flaherty V, and Misstear BD. Using microbiological tracers to assess the impact of winter land use restrictions on the quality of stream headwaters in a small catchment. *Science of the Total Environment* 2016; 541: 949–956. <https://doi.org/10.1016/j.scitotenv.2015.09.071>
- Fouhy F, Ross RP, Fitzgerald G, Stanton C, and Cotter PD. Composition of the early intestinal microbiota: Knowledge, knowledge gaps and the use of high-throughput sequencing to address these gaps. *Gut Microbes* 2012; 3: 203–220. <https://doi.org/10.4161/gmic.20169>
- Fremaux B, Boa T, and Yost CK. Quantitative real-time PCR assays for sensitive detection of Canada goose-specific fecal pollution in water sources. *Applied and Environmental Microbiology* 2010; 76(14): 4886–4889. <https://doi.org/10.1128/AEM.00110-10>
- Gauffin Cano P, Santacruz A, Moya Á, and Sanz Y. *Bacteroides uniformis* CECT 7771 ameliorates metabolic and immunological dysfunction in mice with high-fat-diet induced obesity. *PLoS ONE* 2012; 7(7). <https://doi.org/10.1371/journal.pone.0041079>
- Ge Y, Zhang G, Yang J, Lai XH, Jin D, Lu S, Pu J, Huang Y, Luo X, Zheng H, Zhang X, and Xu J. *Bacteroides luhongzhouii* sp. nov. and *Bacteroides zhangwenhongii* sp. nov., isolated from human faeces. *International Journal of Systematic and Evolutionary Microbiology* 2021; 71: 4772.
- Gibb K, Schobben X, and Christian K. Frogs host faecal bacteria typically associated with humans. *Canadian Journal of Microbiology* 2017; 63(7): 633–637. <https://doi.org/10.1139/cjm-2017-0119>
- Gonzalez RA, Conn KE, Crosswell JR, and Noble RT. Application of empirical predictive modeling using conventional and alternative fecal indicator bacteria in eastern North Carolina waters. *Water Research* 2012; 46(18): 5871–5882. <https://doi.org/10.1016/j.watres.2012.07.050>
- Gordon KV, Brownell M, Wang SY, Lepo JE, Mott J, Nathaniel R, Kilgen M, Hellein K. N, Kennedy E, and Harwood VJ. Relationship of human-associated microbial source tracking markers with enterococci in Gulf of Mexico waters. *Water Research* 2013; 47(3): 996–1004. <https://doi.org/10.1016/j.watres.2012.10.032>
- Gourmelon M, Caprais MP, Mieszkin S, Marti R, Wéry N, Jardé E, Derrien M, Jadas-Hécart A, Communal PY, Jaffrezic A, and Pourcher AM. Development of microbial and chemical MST tools to identify the origin of the faecal pollution in bathing and shellfish harvesting waters in France. *Water Research* 2010; 44(16): 4812–4824. <https://doi.org/10.1016/j.watres.2010.07.061>
- Gourmelon M, Caprais MP, Ségura R, Le Mennec C, Lozach S, Piriou JY, and Rincé A. Evaluation of two library-independent microbial source tracking methods to identify sources of fecal contamination in French estuaries. *Applied and Environmental Microbiology* 2007; 73(15): 4857–4866. <https://doi.org/10.1128/AEM.03003-06>
- Groussin M, Mazel F, Sanders JG, Smillie CS, Lavergne S, Thuiller W, and Alm EJ. Unraveling the processes shaping mammalian gut microbiomes over evolutionary time. *Nature Communications* 2017; 8. <https://doi.org/10.1038/ncomms14319>
- Groussin M, Poyet M, Sistiaga A, Kearney SM, Moniz K, Noel M, Hooker J, Gibbons SM, Segurel L, Froment A, Mohamed RS, Fezeu A, Juimo VA, Lafosse S, Tabe FE, Girard C, Iqaluk D, Nguyen LTT, Shapiro BJ, Lehtimäki J, Ruokolainen L, Kettunen PP, Vatanen T, Sigwazi S, Mabulla A, Dominguez-Rodrigo M, Nartey YA, Agyei-Nkansah A, Duah A, Awuku YA, Valles KA, Asibey SO, Afihene MY, Roberts LR, Plymouth A, Onyekwere CA, Summons RE, Xavier RJ, and Alm EJ. Elevated rates of horizontal gene transfer in the industrialized human microbiome. *Cell* 2021; 184(8): 2053–2067.e18. <https://doi.org/10.1016/j.cell.2021.02.052>
- Guo Y, Kitamoto S, and Kamada N. Microbial adaptation to the healthy and inflamed gut environments. *Gut Microbes* 2020; 12(1): 1–14. <https://doi.org/10.1080/19490976.2020.1857505>
- Hagedorn C, and Xinqiang L. Current and future trends in fecal source tracking and deployment in the Lake Taihu Region of China. *Physics and Chemistry of the Earth* 2011; 36(9–11): 352–359. <https://doi.org/10.1016/j.pce.2010.03.031>
- Hagedorn C, Crozier JB, Mentz KA, Booth AM, Graves AK, Nelson NJ, and Reneau RB. Carbon source utilization profiles as a method to identify sources of faecal pollution in water. *Journal of Applied Microbiology* 2003; 94(5): 792–799. <https://doi.org/10.1046/j.1365-2672.2003.01804.x>
- Hampson D, Crowther J, Bateman I, Kay D, Posen P, Stapleton C, Wyer M, Fezzi C, Jones P, and Tzanopoulos J. Predicting microbial pollution concentrations in UK rivers in response to land use change. *Water Research* 2010; 44(16): 4748–4759. <https://doi.org/10.1016/j.watres.2010.07.062>

- Harwood VJ, Staley C, Badgley BD, Borges K, and Korajkic A. Microbial source tracking markers for detection of fecal contamination in environmental waters: Relationships between pathogens and human health outcomes. *FEMS Microbiology Reviews* 2014; 38(1): 1–40. <https://doi.org/10.1111/1574-6976.12031>
- Hatamoto M, Kaneshige M, Nakamura A, and Yamaguchi T. *Bacteroides luti* sp. nov., an anaerobic, cellulolytic and xylanolytic bacterium isolated from methanogenic sludge. *International Journal of Systematic and Evolutionary Microbiology* 2014; 64(5): 1770–1774. <https://doi.org/10.1099/ijms.0.056630-0>
- Haugland RA, Varma M, Sivaganesan M, Kelty C, Peed L, and Shanks OC. Evaluation of genetic markers from the 16S rRNA gene V2 region for use in quantitative detection of selected Bacteroidales species and human fecal waste by qPCR. *Systematic and Applied Microbiology* 2010; 33(6): 348–357. <https://doi.org/10.1016/j.syapm.2010.06.001>
- He X, Liu P, Zheng G, Chen H, Shi W, Cui Y, Ren H, and Zhang XX. Evaluation of five microbial and four mitochondrial DNA markers for tracking human and pig fecal pollution in freshwater. *Scientific Reports* 2016; 6(May): 1–10. <https://doi.org/10.1038/srep35311>
- He Y, Yuan Q, Mathieu J, Stadler L, Senehi N, Sun R, and Alvarez PJJ. Antibiotic resistance genes from livestock waste: occurrence, dissemination, and treatment. *NPJ Clean Water* 2020; 3(1): 1–11. <https://doi.org/10.1038/s41545-020-0051-0>
- Heaney CD, Myers K, Wing S, Hall D, Baron D, and Stewart JR. Source tracking swine fecal waste in surface water proximal to swine concentrated animal feeding operations. *Science of the Total Environment* 2015; 511: 676–683. <https://doi.org/10.1016/j.scitotenv.2014.12.062>
- Holdeman LV, and Moore W. New genus, *Coprococcus*, twelve new species, and emended descriptions of four previously described species of bacteria from human feces. *International Journal of Systematic Bacteriology* 1974; 24(2): 260–277.
- Hong PY, Wu JH, and Liu WT. Relative abundance of *Bacteroides* spp. in stools and wastewaters as determined by hierarchical oligonucleotide primer extension. *Applied and Environmental Microbiology* 2008; 74(9): 2882–2893. <https://doi.org/10.1128/AEM.02568-07>
- Huddleston JR. Horizontal gene transfer in the human gastrointestinal tract: Potential spread of antibiotic resistance genes. *Infection and Drug Resistance* 2014; 7: 167–176. <https://doi.org/10.2147/IDR.S48820>
- Husain F, Tang K, Veeranagouda Y, Boente R, Patrick S, Blakely G, and Wexler HM. Novel large-scale chromosomal transfer in *Bacteroides fragilis* contributes to its pan-genome and rapid environmental adaptation. *Microbial Genomics* 2017; 3(11). <https://doi.org/10.1099/mgen.0.000136>
- Irisawa T, Saputra S, Kitahara M, Sakamoto M, Sulistiani, Yulineri T, Dinoto A, and Ohkuma M. *Bacteroides caecicola* sp. nov. and *Bacteroides gallinaceum* sp. nov., isolated from the caecum of an Indonesian chicken. *International Journal of Systematic and Evolutionary Microbiology* 2016; 66(3): 1431–1437. <https://doi.org/10.1099/ijsem.0.000899>
- Jamwal P, Mittal AK, and Mouchel JM. Point and non-point microbial source pollution: A case study of Delhi. *Physics and Chemistry of the Earth* 2011; 36(12): 490–499. <https://doi.org/10.1016/j.pce.2008.09.005>
- Jardé E, Jeanneau L, Harrault L, Quenot E, Solecki O, Petitjean P, Lozach S, Chev e J, and Gourmelon M. Application of a microbial source tracking based on bacterial and chemical markers in headwater and coastal catchments. *Science of the Total Environment* 2018; 610–611: 55–63. <https://doi.org/10.1016/j.scitotenv.2017.07.235>
- Jenkins MW, Tiwari S, Lorente M, Gichaba CM, and Wuertz S. Identifying human and livestock sources of fecal contamination in Kenya with host specific Bacteroidales assays. *Water Research* 2009; 43(19): 4956–4966. <https://doi.org/10.1016/j.watres.2009.07.028>
- Jensen NS, and Canale-Parola E. *Bacteroides pectinophilus* sp. nov. and *Bacteroides galacturonicus* sp. nov.: Two pectinolytic bacteria from the human intestinal tract. *Applied and Environmental Microbiology* 1986; 52(4): 880–887. <https://doi.org/10.1128/aem.52.4.880-887.1986>
- Johnson JL, Moore WEC, and Moore, LVH. *Bacteroides caccae* sp. nov., *Bacteroides merdae* sp. nov., and *Bacteroides stercoris* sp. nov. isolated from human feces. *International Journal of Systematic Bacteriology* 1986; 1: 499–501.
- Kabiri L, Alum A, Rock C, McLain JE, and Abbaszadegan M. Isolation of *Bacteroides* from fish and human fecal samples for identification of unique molecular markers. *Canadian Journal of Microbiology* 2013; 59(12): 771–777. <https://doi.org/10.1139/cjm-2013-0518>
- Kierzkowska M, Majewska A, and Mlynarczyk G. Trends and impact in antimicrobial resistance among *Bacteroides* and *Parabacteroides* species in 2007–2012 compared to 2013–2017. *Microbial Drug Resistance* 2020; 26(12): 1452–1457. <https://doi.org/10.1089/mdr.2019.0462>
- Kildare BJ, Leutenegger CM, Mcswain BS, Bambic DG, Rajal VB, and Wuertz S. 16S rRNA-based assays for quantitative detection of Bacteroidales: A Bayesian approach. *Water* 2007; 41: 3701–3715. <https://doi.org/10.1016/j.watres.2007.06.037>
- Kim MS, Roh SW, and Bae JW. *Bacteroides faecis* sp. nov., isolated from human faeces. *International Journal of Systematic and Evolutionary Microbiology* 2010; 60(11): 2572–2576. <https://doi.org/10.1099/ijms.0.020024-0>
- Kirschner AKT, Reischer GH, Jakwerth S, Savio D, Ixenmaier S, Toth E, Sommer R, Mach RL, Linke R, Eiler A, Kolarevic S, and Farnleitner AH. Multiparametric monitoring of microbial faecal pollution reveals the dominance of human contamination along the whole Danube River. *Water Research* 2017; 124: 543–555. <https://doi.org/10.1016/j.watres.2017.07.052>
- Kitahara M, Sakamoto M, Tsuchida S, Kawasumi K, Amao H, Benno Y, and Ohkuma, M. *Bacteroides stercorisoris* sp. nov. and *Bacteroides faecichinchillae* sp. nov., isolated from chinchilla (*Chinchilla lanigera*) faeces. *International Journal of Systematic and Evolutionary Microbiology* 2012; 62(5): 1145–1150. <https://doi.org/10.1099/ijms.0.032706-0>
- Kitahara M, Tsuchida S, Kawasumi K, Amao H, Sakamoto M, Benno Y, and Ohkuma M. *Bacteroides chinchillae* sp. nov. and *Bacteroides rodentium* sp. nov., isolated from chinchilla (*Chinchilla lanigera*) faeces. *International Journal of Systematic and Evolutionary Microbiology* 2011; 61(4): 877–881. <https://doi.org/10.1099/ijms.0.024026-0>

- Knappett PSK, Escamilla V, Layton A, McKay LD, Emch M, Williams DE, Huq R, Alam, J, Farhana L, Mailloux BJ, Ferguson A, Saylor GS, Ahmed KM, and van Geen A. Impact of population and latrines on fecal contamination of ponds in rural Bangladesh. *Science of the Total Environment* 2011; 409(17): 3174–3182. <https://doi.org/10.1016/j.scitotenv.2011.04.043>
- Kobayashi A, Sano D, Hatori J, Ishii S, and Okabe S. Chicken- and duck-associated *Bacteroides-Prevotella* genetic markers for detecting fecal contamination in environmental water. *Applied Microbiology and Biotechnology* 2013; 97(16): 7427–7437. <https://doi.org/10.1007/s00253-012-4469-2>
- Lagkouvardos I, Pukall R, Abt B, Foessel BU, Meier-Kolthoff JP, Kumar N, Bresciani A, Martínez I, Just S, Ziegler C, Brugiroux S, Garzetti D, Wenning M, Bui TPN, Wang J, Hugenholtz F, Plugge CM, Peterson DA, Hornef MW, Baines JF, Smidt H, Walter J, Kristiansen K, Nielsen HB, Haler D, Overmann J, Stecher B, and Clavel T. The Mouse Intestinal Bacterial Collection (miBC) provides host-specific insight into cultured diversity and functional potential of the gut microbiota. *Nature Microbiology* 2016; 1(October): 1–16. <https://doi.org/10.1038/nmicrobiol.2016.131>
- Lan PTN, Sakamoto M, Sakata S, and Benno Y. *Bacteriodes barnesiae* sp. nov., *Bacteroides salanitronis* sp. nov. and *Bacteroides gallinarum* sp. nov., isolated from chicken caecum. *International Journal of Systematic and Evolutionary Microbiology* 2006; 56(12): 2853–2859. <https://doi.org/10.1099/ijs.0.64517-0>
- Landuyt AE, Klocke BJ, Duck LW, Kemp KM, Muir RQ, Jennings MS, Blum SI, Tse HM, Lee G, Morrow CD, Elson CO, and Maynard CL. ICOS ligand and IL-10 synergize to promote host-microbiota mutualism. *Proceedings of the National Academy of Sciences of the United States of America* 2021; 118(13): 1–11. <https://doi.org/10.1073/pnas.2018278118>
- Layton A, McKay L, Williams D, Garrett V, Gentry R, and Saylor G. Development of *Bacteroides* 16S rRNA gene TaqMan-based real-time PCR assays for estimation of total, human, and bovine fecal pollution in water. *Applied and Environmental Microbiology* 2006; 72(6): 4214–4224. <https://doi.org/10.1128/AEM.01036-05>
- Lee CS, and Lee J. Application of host-specific source-tracking tools for rapid identification of fecal contamination in fresh produce by humans and livestock. *Journal of the Science of Food and Agriculture* 2012; 93(5): 1089–1096. <https://doi.org/10.1002/jsfa.5855>
- Lee JE, Lee H, Cho YH, Hur HG, and Ko GP. F+ RNA coliphage-based microbial source tracking in water resources of South Korea. *Science of the Total Environment*, 2011; 412–413: 127–131. <https://doi.org/10.1016/j.scitotenv.2011.09.061>
- Ley RE, Lozupone, CA, Hamady M, Knight R, and Gordon JJ. Worlds within worlds: Evolution of the vertebrate gut microbiota. *Nature Reviews Microbiology* 2008; 6(10): 776–788. <https://doi.org/10.1038/nrmicro1978>
- Li E, Saleem F, Edge TA, and Schellhorn HE. Biological indicators for fecal pollution detection and source tracking: A review. *Processes* 2021a; 9(11). <https://doi.org/10.3390/pr9112058>
- Li X, Kang Y, Huang Y, Xiao Y, Song L, Lu S, and Ren Z. A strain of *Bacteroides thetaiotaomicron* attenuates colonization of *Clostridioides difficile* and affects intestinal microbiota and bile acids profile in a mouse model. *Biomedicine and Pharmacotherapy* 2021b; 137: 11290. <https://doi.org/10.1016/j.biopha.2021.111290>
- Ma T, McAllister TA, and Guan LL. A review of the resistome within the digestive tract of livestock. *Journal of Animal Science and Biotechnology* 2021; 12(1), 1–20. <https://doi.org/10.1186/s40104-021-00643-6>
- Maceda-Veiga A, Mac Nally R, and de Sostoa A. The presence of non-native species is not associated with native fish sensitivity to water pollution in greatly hydrologically altered rivers. *Science of the Total Environment* 2017; 607–608: 549–557. <https://doi.org/10.1016/j.scitotenv.2017.07.010>
- Majewska A, Kierzkowska M, and Kawecki D. What we actually know about the pathogenicity of *Bacteroides pyogenes*. *Medical Microbiology and Immunology* 2021; 210(2–3): 157–163. <https://doi.org/10.1007/s00430-021-00709-2>
- Markand S, Bachoon DS, Gentit L, Sherchan S, and Gates K. Evaluation of physical, chemical and microbiological parameters of water quality in the Harris Neck estuarine marshes along the Georgia coast. *Marine Pollution Bulletin* 2011; 62(1): 178–181. <https://doi.org/10.1016/j.marpolbul.2010.11.010>
- Marti R, Gannon VPJ, Jokinen C, Lanthier M, Lapen DR, Neumann NF, Ruecker NJ, Scott A, Wilkes G, Zhang Y, and Topp E. Quantitative multi-year elucidation of fecal sources of waterborne pathogen contamination in the South Nation River basin using Bacteroidales microbial source tracking markers. *Water Research* 2013; 47(7): 2315–2324. <https://doi.org/10.1016/j.watres.2013.02.009>
- Marti R, Zhang Y, Lapen DR, and Topp E. Development and validation of a microbial source tracking marker for the detection of fecal pollution by muskrats. *Journal of Microbiological Methods* 2011; 87(1): 82–88. <https://doi.org/10.1016/j.mimet.2011.07.013>
- Matsuoka T, Shimizu T, Minagawa T, Hiranuma W, Takeda M, Kakuta R, and Kawamoto S. First case of an invasive *Bacteroides dorei* infection detected in a patient with a mycotic aortic aneurysm—raising a rebellion of major indigenous bacteria in humans: a case report and review. *BMC Infectious Diseases* 2021; 21(1): 1–6. <https://doi.org/10.1186/s12879-021-06345-8>
- Mattioli MC, Sassoubre LM, Russell TL, and Boehm AB. Decay of sewage-sourced microbial source tracking markers and fecal indicator bacteria in marine waters. *Water Research* 2017; 108: 106–114. <https://doi.org/10.1016/j.watres.2016.10.066>
- Mauffret A, Caprais MP, and Gourmelon M. Relevance of Bacteroidales and F-specific RNA bacteriophages for efficient fecal contamination tracking at the level of a catchment in France. *Applied and Environmental Microbiology* 2012; 78(15): 5143–5152. <https://doi.org/10.1128/AEM.00315-12>
- Mauffret A, Mieszkin S, Morizur M, Alfiansah Y, Lozach S, and Gourmelon M. Recent innovation in microbial source tracking using bacterial real-time PCR markers in shellfish. *Marine*

- Pollution Bulletin 2013; 68(1–2): 21–29. <https://doi.org/10.1016/j.marpolbul.2012.12.029>
- Mayer RE, Bo S, Egle L, Reischer GH, Schade M, Fernandez-Cassi X, Fuchs W, Mach RL, Lindner G, Kirschner A, Gaisbauer M, Piringer H, Blaschke AP, Girones R, Zessner M, Sommer R, and Farnleitner AH. Occurrence of human-associated Bacteroidetes genetic source tracking markers in raw and treated wastewater of municipal and domestic origin and comparison to standard and alternative indicators of faecal pollution. *Water Research* 2016; 90: 256–276. <https://doi.org/10.1016/j.watres.2015.12.031>
- Mazmanian SK, and Kasper DL. The love-hate relationship between bacterial polysaccharides and the host immune system. *Nature Reviews Immunology* 2006; 6(11): 849–858. <https://doi.org/10.1038/nri1956>
- Meays CL, Broersma K, Nordin R, and Mazumder A. Source tracking fecal bacteria in water: a critical review of current methods. *Journal of Environmental Management* 2004; 73: 71–79. <https://doi.org/10.1016/j.jenvman.2004.06.001>
- Mieszkin S, Furet JP, Corthier G, and Gourmelon M. Estimation of pig fecal contamination in a river catchment by real-time PCR using two pig-specific Bacteroidales 16S rRNA genetic markers. *Applied and Environmental Microbiology* 2009; 75(10): 3045–3054. <https://doi.org/10.1128/AEM.02343-08>
- Mieszkin S, Yala JF, Joubrel R, and Gourmelon M. Phylogenetic analysis of Bacteroidales 16S rRNA gene sequences from human and animal effluents and assessment of ruminant faecal pollution by real-time PCR. *Journal of Applied Microbiology* 2010; 108(3): 974–984. <https://doi.org/10.1111/j.1365-2672.2009.04499.x>
- Milani C, Duranti S, Bottacini F, Casey E, Turroni F, Mahony J, Belzer C, Delgado Palacio S, Arboleya Montes S, Mancabelli L, Lugli GA, Rodriguez JM, Bode L, de Vos W, Gueimonde M, Margolles A, van Sinderen D, and Ventura M. The First microbial colonizers of the human gut: Composition, activities, and health implications of the infant gut microbiota. *Microbiology and Molecular Biology Reviews* 2017; 81(4): 1–67. <https://doi.org/10.1128/membr.00036-17>
- Miyamoto Y, and Itoh K. *Bacteroides acidifaciens* sp. nov., isolated from the caecum of mice. *International Journal of Systematic and Evolutionary Microbiology* 2000; 50(1): 145–148. <https://doi.org/10.1099/00207713-50-1-145>
- Mori K, and Kamagata Y. The challenges of studying the anaerobic microbial world. *Microbes and Environments* 2014; 29(4): 335–337. <https://doi.org/10.1264/jsme2.ME2904rh>
- Niestępski S, Harnisz M, Korzeniewska E, and Osińska A. Isolation of anaerobic bacteria of the *Bacteroides fragilis* group from environmental samples. *E3S Web of Conferences* 2019; 100. <https://doi.org/10.1051/e3sconf/201910000058>
- Nishiyama T, Ueki A, Kaku N, Watanabe K, and Ueki K. *Bacteroides graminisolvens* sp. nov., a xylanolytic anaerobe isolated from a methanogenic reactor treating cattle waste. *International Journal of Systematic and Evolutionary Microbiology* 2009; 59(8): 1901–1907. <https://doi.org/10.1099/ijs.0.008268-0>
- Noble RT, Griffith JF, Blackwood AD, Fuhrman JA, Gregory JB, Hernandez X, Liang X, Bera AA, and Schiff, K. Multitiered approach using quantitative PCR to track sources of fecal pollution affecting Santa Monica Bay, California. *Applied and Environmental Microbiology* 2006; 72(2): 1604–1612. <https://doi.org/10.1128/AEM.72.2.1604-1612.2006>
- Noorhosseini SA, Allahyari MS, Damalas CA, and Moghaddam SS. Public environmental awareness of water pollution from urban growth: The case of Zarjub and Goharrud rivers in Rasht, Iran. *Science of the Total Environment* 2017; 599–600: 2019–2025. <https://doi.org/10.1016/j.scitotenv.2017.05.128>
- Nshimiyimana JP, Ekklesia E, Shanahan P, Chua LHC, and Thompson JR. Distribution and abundance of human specific *Bacteroides* and relation to traditional indicators in an urban tropical catchment. *Journal of Applied Microbiology* 2014; 116(5): 1369–1383. <https://doi.org/10.1111/jam.12455>
- O’Toole PW, Marchesi JR, and Hill C. Next-generation probiotics: The spectrum from probiotics to live biotherapeutics. *Nature Microbiology* 2017; 2(April): 1–6. <https://doi.org/10.1038/nmicrobiol.2017.57>
- Ochoa-Repáraz J, Mielcarz DW, Ditrio LE, Burroughs AR, Begum-Haque S, Dasgupta S, Kasper DL, and Kasper LH. Central nervous system demyelinating disease protection by the human commensal *Bacteroides fragilis* depends on polysaccharide A expression. *Journal of Immunology* 2010; 185(7): 4101–4108. <https://doi.org/10.4049/jimmunol.1001443>
- Odagiri M, Schriewer A, Hanley K, Wuertz S, Misra PR, Panigrahi P, and Jenkins MW. Validation of Bacteroidales quantitative PCR assays targeting human and animal fecal contamination in the public and domestic domains in India. *Science of the Total Environment* 2015; 502: 462–470. <https://doi.org/10.1016/j.scitotenv.2014.09.040>
- Ogunkunle CO, Mustapha K, Oyedjeji S, and Fatoba PO. Assessment of metallic pollution status of surface water and aquatic macrophytes of earthen dams in Ilorin, north-central of Nigeria as indicators of environmental health. *Journal of King Saud University – Science* 2016; 28(4): 324–331. <https://doi.org/10.1016/j.jksus.2015.11.005>
- Ohad S, Vaizel-Ohayon D, Rom M, Guttman J, Berger D, Kravitz V, Pilo S, Huberman Z, Kashi Y, and Rorman E. Microbial source tracking in adjacent karst springs. *Applied and Environmental Microbiology* 2015; 81(15): 5037–5047. <https://doi.org/10.1128/AEM.00855-15>
- Okabe S, Okayama N, Savichtcheva O, and Ito T. Quantification of host-specific *Bacteroides-Prevotella* 16S rRNA genetic markers for assessment of fecal pollution in freshwater. *Applied Microbiology and Biotechnology* 2007; 74(4): 890–901. <https://doi.org/10.1007/s00253-006-0714-x>
- Oun A, Yin Z, Munir M, and Xagorarakis I. Microbial pollution characterization of water and sediment at two beaches in Saginaw Bay, Michigan. *Journal of Great Lakes Research* 2017; 43(3): 64–72. <https://doi.org/10.1016/j.jglr.2017.01.014>
- Paruch L, Paruch AM, Eiken HG, and Sørheim R. Faecal pollution affects abundance and diversity of aquatic microbial community in anthropo-zoogenically influenced lotic ecosystems. *Scientific Reports* 2019; 9(1): 1–13. <https://doi.org/10.1038/s41598-019-56058-x>
- Popkin BM, D’Anci KE, and Rosenberg IH. Water, hydration, and health. *Nutrition Reviews* 2010; 68(8), 439–458. <https://doi.org/10.1111/j.1753-4887.2010.00304.x>

- Ram JL, Thompson B, Turner C, Nechvtal JM, Sheehan H, and Bobrin J. Identification of pets and raccoons as sources of bacterial contamination of urban storm sewers using a sequence-based bacterial source tracking method. *Water Research* 2007; 41: 3605–3614. <https://doi.org/10.1016/j.watres.2007.04.013>
- Ravaliya K, Gentry-Shields J, Garcia S, Heredia N, de Aceituno AF, Bartz FE, Leon J. S, and Jaykus LA. Use of Bacteroidales microbial source tracking to monitor fecal contamination in fresh produce production. *Applied and Environmental Microbiology* 2014; 80(2): 612–617. <https://doi.org/10.1128/AEM.02891-13>
- Reed KD, Meece JK, Henkel JS, and Shukla SK. Birds, migration and emerging zoonoses: West Nile virus, Lyme disease, influenza A and enteropathogens. *Clinical Medicine & Research* 2003; 1(1): 5–12. <https://doi.org/10.3121/cmr.1.1.5>
- Reischer GH, Kasper DC, Steinborn R, Farnleitner AH, and Mach RL. A quantitative real-time PCR assay for the highly sensitive and specific detection of human faecal influence in spring water from a large alpine catchment area. *Letters in Applied Microbiology* 2007; 44(4): 351–356. <https://doi.org/10.1111/j.1472-765X.2006.02094.x>
- Reischer GH, Kasper DC, Steinborn R, Mach RL, and Farnleitner AH. Quantitative PCR method for sensitive detection of ruminant fecal pollution in freshwater and evaluation of this method in alpine karstic regions. *Applied and Environmental Microbiology* 2006 72(8): 5610–5614. <https://doi.org/10.1128/AEM.00364-06>
- Ridley CM, Jamieson RC, Truelstrup Hansen L, Yost CK, and Bezanson GS. Baseline and storm event monitoring of Bacteroidales marker concentrations and enteric pathogen presence in a rural Canadian watershed. *Water Research* 2014; 60: 278–288. <https://doi.org/10.1016/j.watres.2014.04.039>
- Rivera B, and Rock C. Microbial source tracking: watershed characterization and source identification. *Arizona Cooperative Extension* 2011; 1–8.
- Robert C, Chassard C, Lawson PA, and Bernalier-Donadille A. *Bacteroides cellulosilyticus* sp. nov., a cellulolytic bacterium from the human gut microbial community. *International Journal of Systematic and Evolutionary Microbiology* 2007; 57(7): 1516–1520. <https://doi.org/10.1099/ijs.0.64998-0>
- Ruzol C, Banzon-Cabanilla D, Ancog R, and Peralta E. Understanding water pollution management: Evidence and insights from incorporating cultural theory in social network analysis. *Global Environmental Change* 2017; 45(July): 183–193. <https://doi.org/10.1016/j.gloenvcha.2017.06.009>
- Sadowski C, and Bowman J. Historical surveys reveal a long-term decline in muskrat populations. *Ecology and Evolution* 2021; 11(12): 7557–7568. <https://doi.org/10.1002/ece3.7588>
- Sakamoto M, and Ohkuma M. *Bacteroides reticulotermitis* sp. nov., isolated from the gut of a subterranean termite (*Reticulitermes speratus*). *International Journal of Systematic and Evolutionary Microbiology* 2013; 63(2): 691–695. <https://doi.org/10.1099/ijs.0.040931-0>
- Saputra S, Irisawa T, Sakamoto M, Kitahara M, Sulistiani, Yulinery T, Ohkuma M, and Dinoto A. *Bacteroides caecigallinarum* sp. Nov., isolated from caecum of an Indonesian chicken. *International Journal of Systematic and Evolutionary Microbiology* 2015; 65(12): 4341–4346. <https://doi.org/10.1099/ijsem.0.000573>
- Sauer EP, VandeWalle JL, Bootsma MJ, and McLellan SL. Detection of the human specific *Bacteroides* genetic marker provides evidence of widespread sewage contamination of stormwater in the urban environment. *Water Research* 2011; 45(14): 4081–4091. <https://doi.org/10.1016/j.watres.2011.04.049>
- Savichtcheva O, and Okabe S. Alternative indicators of fecal pollution: Relations with pathogens and conventional indicators, current methodologies for direct pathogen monitoring and future application perspectives. *Water Research* 2006; 40(13): 2463–2476. <https://doi.org/10.1016/j.watres.2006.04.040>
- Schriewer A, Miller WA, Byrne BA, Miller MA, Oates S, Conrad PA, Hardin D, Yang HH, Chouicha N, Melli A, Jessup D, Dominik C, and Wuertz S. Presence of Bacteroidales as a predictor of pathogens in surface waters of the central California coast. *Applied and Environmental Microbiology* 2010; 76(17), 5802–5814. <https://doi.org/10.1128/AEM.00635-10>
- Shanks OC, Kelty CA, Sivaganesan M, Varma M, and Haugland RA. Quantitative PCR for genetic markers of human fecal pollution. *Applied and Environmental Microbiology* 2009; 75(17): 5507–5513. <https://doi.org/10.1128/AEM.00305-09>
- Shin Y, Park SJ, Paek J, Kim JS, Rhee MS, Kim H, Kook JK, and Chang YH. *Bacteroides koreensis* sp. nov. and *Bacteroides kribbi* sp. nov., two new members of the genus *Bacteroides*. *International Journal of Systematic and Evolutionary Microbiology* 2017; 67(11): 4352–4357. <https://doi.org/10.1099/ijsem.0.002226>
- Smalley D, Rocha ER, and Smith CJ. Aerobic-type ribonucleotide reductase in the anaerobe *Bacteroides fragilis*. *Journal of Bacteriology* 2002; 184(4): 895–903. <https://doi.org/10.1128/jb.184.4.895-903.2002>
- Sóki J, Wybo I, Hajdú E, Toprak NU, Jeverica S, Stingu CS, Tierney D, Perry JD, Matuz M, Urbán E, and Nagy E. A Europe-wide assessment of antibiotic resistance rates in *Bacteroides* and *Parabacteroides* isolates from intestinal microbiota of healthy subjects. *Anaerobe* 2020; 62. <https://doi.org/10.1016/j.anaerobe.2020.102182>
- Somnark P, Chyerochana N, Kongprajug A, Mongkolsuk S, and Sirikanchana K. PCR data and comparative performance of Bacteroidales microbial source tracking genetic markers. *Data in Brief* 2018a; 19: 156–169. <https://doi.org/10.1016/j.dib.2018.04.129>
- Somnark P, Chyerochana N, Mongkolsuk S, and Sirikanchana K. Performance evaluation of Bacteroidales genetic markers for human and animal microbial source tracking in tropical agricultural watersheds. *Environmental Pollution* 2018b; 236: 100–110. <https://doi.org/10.1016/j.envpol.2018.01.052>
- Song YL, Liu CX, Mctague M, and Finegold SM. Isolated from clinical specimens of human intestinal origin. *Society* 2004; 42(12): 5565–5570. <https://doi.org/10.1128/JCM.42.12.5565>
- Staley C, Reckhow KH, Lukasik J, and Harwood VJ. Assessment of sources of human pathogens and fecal contamination in a Florida freshwater lake. *Water Research*

- 2012; 46(17): 5799–5812.  
<https://doi.org/10.1016/j.watres.2012.08.012>
- Stapleton CM, Kay D, Wyer MD, Davies C, Watkins J, Kay C, McDonald AT, Porter J, and Gawler A. Evaluating the operational utility of a Bacteroidales quantitative PCR-based MST approach in determining the source of faecal indicator organisms at a UK bathing water. *Water Research* 2009; 43(19): 4888–4899.  
<https://doi.org/10.1016/j.watres.2009.09.015>
- Stea EC, Truelstrup Hansen L, Jamieson RC, and Yost CK. Fecal contamination in the surface waters of a rural- and an urban-source watershed. *Journal of Environment Quality* 2015; 44(5): 1556–1567.  
<https://doi.org/10.2134/jeq2014.11.0459>
- Strandwitz P, Kim KH, Terekhova D, Liu JK, Sharma A, Levering J, McDonald D, Dietrich D, Ramadhar TR, Lekbua A, Mroue N, Liston C, Stewart EJ, Dubin MJ, Zengler K, Knight R, Gilbert JA, Clardy J, and Lewis K. GABA-modulating bacteria of the human gut microbiota. *Nature Microbiology* 2019; 4(3): 396–403.  
<https://doi.org/10.1038/s41564-018-0307-3>
- Symonds EM, Young S, Verbyla ME, McQuaig-Ulrich SM, Ross E, Jimenez JA, Harwood VJ, and Breitbart M. Microbial source tracking in shellfish harvesting waters in the Gulf of Nicoya, Costa Rica. *Water Research* 2017; 111: 177–184.  
<https://doi.org/10.1016/j.watres.2017.01.004>
- Tamana SK, Tun HM, Konya T, Chari RS, Field CJ, Guttman DS, Becker AB, Moraes TJ, Turvey SE, Subbarao P, Sears MR, Pei J, Scott JA, Mandhane PJ, and Kozyrskyj AL. *Bacteroides*-dominant gut microbiome of late infancy is associated with enhanced neurodevelopment. *Gut Microbes* 2021; 13(1): 1–17.  
<https://doi.org/10.1080/19490976.2021.1930875>
- Tambalo DD, Boa T, Liljebjelke K, and Yost CK. Evaluation of two quantitative PCR assays using Bacteroidales and mitochondrial DNA markers for tracking dog fecal contamination in waterbodies. *Journal of Microbiological Methods* 2012; 91(3): 459–467.  
<https://doi.org/10.1016/j.mimet.2012.09.029>
- Teixeira AF, Kühnel W, Vives P, and Wedel T. Functional morphology of unguiculiiform papillae of the reticular groove in the ruminant stomach. *Annals of Anatomy* 2009; 191(5): 469–476.  
<https://doi.org/10.1016/j.aanat.2009.06.005>
- Thornton RF, Murphy EC, Kagawa TF, Otoole PW, and Cooney JC. The effect of environmental conditions on expression of *Bacteroides fragilis* and *Bacteroides thetaiotaomicron* C10 protease genes. *BMC Microbiology* 2012; 12.  
<https://doi.org/10.1186/1471-2180-12-190>
- Troy EB, and Kasper DL. Beneficial effects of *Bacteroides fragilis* polysaccharides on the immune system. *Frontiers in Bioscience* 2010; 28(2): 25–34.  
<https://doi.org/10.1080/17843286.1973.11716863>
- Turrone F, Milani C, Duranti S, Lugli GA, Bernasconi S, Margolles A, Di Pierro F, Van Sinderen D, and Ventura M. The infant gut microbiome as a microbial organ influencing host well-being. *Italian Journal of Pediatrics* 2020; 46(1): 1–13.  
<https://doi.org/10.1186/s13052-020-0781-0>
- Ueki A, Abe K, Kaku N, Watanabe K, and Ueki K. *Bacteroides propionicifaciens* sp. nov., isolated from rice-straw residue in a methanogenic reactor treating waste from cattle farms. *International Journal of Systematic and Evolutionary Microbiology* 2008; 58(2): 346–352.  
<https://doi.org/10.1099/ijs.0.65486-0>
- Ulsemer P, Toutounian K, Schmidt J, Karsten U, and Goletz S. Preliminary safety evaluation of a new *Bacteroides xylanisolvans* isolate. *Applied and Environmental Microbiology* 2012; 78(2), 528–535.  
<https://doi.org/10.1128/AEM.06641-11>
- Vadde KK, Feng Q, Wang J, McCarthy AJ, and Sekar R. Next-generation sequencing reveals fecal contamination and potentially pathogenic bacteria in a major inflow river of Taihu Lake. *Environmental Pollution* 2019; 254: 113108.  
<https://doi.org/10.1016/j.envpol.2019.113108>
- Varki A. Biological roles of glycans. *Glycobiology* 2017; 27(1): 3–49.  
<https://doi.org/10.1093/glycob/cww086>
- Villemur R, Imbeau M, Vuong MN, Masson L, and Payment P. An environmental survey of surface waters using mitochondrial DNA from human, bovine and porcine origin as fecal source tracking markers. *Water Research* 2015; 69: 143–153.  
<https://doi.org/10.1016/j.watres.2014.10.063>
- Vrzal J, Vuković-Gačić B, Kolarević S, Gačić Z, Kračun-Kolarević M, Kostić J, Aborigiba M, Farnleitner A, Reischer G, Linke R, Paunović M, and Ogrinc N. Determination of the sources of nitrate and the microbiological sources of pollution in the Sava River Basin. *Science of the Total Environment* 2016; 573: 1460–1471.  
<https://doi.org/10.1016/j.scitotenv.2016.07.213>
- Wang M, Li M, Wu S, Lebrilla CB, Chapkin RS, Ivanov I, and Donovan SM. Fecal microbiota composition of breast-fed infants is correlated with human milk oligosaccharides consumed. *Journal of Pediatric Gastroenterology and Nutrition* 2015; 60(6): 825–833.  
<https://doi.org/10.1097/MPG.0000000000000752>
- Watanabe Y, Nagai F, Morotomi M, Sakon H, and Tanaka R. *Bacteroides clarus* sp. nov., *Bacteroides fluxus* sp. nov. and *Bacteroides oleiciplenus* sp. nov., isolated from human faeces. *International Journal of Systematic and Evolutionary Microbiology* 2010; 60(8): 1864–1869.  
<https://doi.org/10.1099/ijs.0.015107-0>
- Weiner LM, Webb AK, Limbago B, Dudeck MA, Patel J, Kallen AJ, Edwards JR, and Sievert DM. Antimicrobial-resistant pathogens associated with healthcare-associated infections: summary of data reported to the National Healthcare Safety Network at the Centers for Disease Control and Prevention, 2011–2014. *Infection Control and Hospital Epidemiology* 2016; 37(11): 1288–1301.  
<https://doi.org/10.1017/ice.2016.174>
- Wexler AG, and Goodman AL. An insider’s perspective: *Bacteroides* as a window into the microbiome. *Nature Microbiology* 2017; 2(April): 1–11.  
<https://doi.org/10.1038/nmicrobiol.2017.26>
- Wexler HM. *Bacteroides*: The good, the bad, and the nitty-gritty. *Clinical Microbiology Reviews* 2007; 20(4): 593–621.  
<https://doi.org/10.1128/CMR.00008-07>
- Whitehead TR, Cotta MA, Collins MD, Falsen E, and Lawson PA. *Bacteroides coprosuis* sp. nov., isolated from swine-manure storage pits. *International Journal of Systematic and*



- Evolutionary Microbiology 2005; 55(6): 2515–2518. <https://doi.org/10.1099/ijs.0.63869-0>
- Wiggins BA. Discriminant analysis of antibiotic resistance patterns in fecal streptococci, a method to differentiate human and animal sources of fecal pollution in natural waters. Applied and Environmental Microbiology 1996; 62(11): 3997–4002.
- Wong K, Fong T, Bibby K, and Molina M. Application of enteric viruses for fecal pollution source tracking in environmental waters. Environment International 2012; 45: 151–164. <https://doi.org/10.1016/j.envint.2012.02.009>
- Woodall CJ. Waterborne diseases - What are the primary killers? Desalination 2009; 248(1–3): 616–621. <https://doi.org/10.1016/j.desal.2008.05.110>
- Xue J, Lin S, Lamar F, Lamori, JG, and Sherchan S. Assessment of fecal pollution in Lake Pontchartrain, Louisiana. Marine Pollution Bulletin 2018; 129(2): 655–663. <https://doi.org/10.1016/j.marpolbul.2017.10.053>
- Yan, Zhang, W, Zhang Z, Liu Y, and Deng C. Assessment of water quality and identification of polluted risky regions based on field observations & GIS in the Honghe River Watershed, China. PLOS ONE 2015; 1–13. <https://doi.org/10.1371/journal.pone.0119130>
- Yanagibashi T, Hosono A, Oyama A, Tsuda M, Suzuki A, Hachimura S, Takahashi Y, Momose Y, Itoh K, Hirayama K, Takahashi K, & Kaminogawa S. IgA production in the large intestine is modulated by a different mechanism than in the small intestine: *Bacteroides acidifaciens* promotes IgA production in the large intestine by inducing germinal center formation and increasing the number of IgA+ B cell. Immunobiology 2013; 218(4): 645–651. <https://doi.org/10.1016/j.imbio.2012.07.033>
- Yu SY, Kim JS, Oh BS, Park SH, Kang SW, Park JE, Choi SH, Han K Il, Lee KC, Eom MK, Suh MK, Lee DH, Yoon H, Kim BY, Yang SJ, Lee JS, and Lee JH. *Bacteroides faecalis* sp. nov., isolated from human faeces. International Journal of Systematic and Evolutionary Microbiology 2019; 69(12): 3824–3829. <https://doi.org/10.1099/ijsem.0.003690>
- Zafar H, and Saier MH. Comparative genomics of transport proteins in seven *Bacteroides* species. PLoS ONE 2018; 13(12): 1–25. <https://doi.org/10.1371/journal.pone.0208151>
- Zafar H, and Saier MH. Gut *Bacteroides* species in health and disease. Gut Microbes 2021; 13(1), 1–20. <https://doi.org/10.1080/19490976.2020.1848158>
- Zamani S, Taslimi R, Sarabi A, Jasemi S, Sechi LA, and Feizabadi MM. Enterotoxigenic *Bacteroides fragilis*: A possible etiological candidate for bacterially-induced colorectal precancerous and cancerous lesions. Frontiers in Cellular and Infection Microbiology 2020; 9(1): 1–7. <https://doi.org/10.3389/fcimb.2019.00449>
- Zhang Q, Eichmiller JJ, Staley C, Sadowsky MJ, and Ishii S. Correlations between pathogen concentration and fecal indicator marker genes in beach environments. Science of the Total Environment 2016; 573: 826–830. <https://doi.org/10.1016/j.scitotenv.2016.08.122>
- Zheng G, and Shen Z. Host-specific genetic markers of fecal bacteria for fecal source tracking in food and water. Journal of Food: Microbiology, Safety & Hygiene 2018; 3(01): 1–8. <https://doi.org/10.4172/2476-2059.1000135>
- Zhong H, Penders J, Shi Z, Ren H, Cai K, Fang C, Ding Q, Thijs C, Blaak EE, Stehouwer CDA, Xu X, Yang H, Wang J, Wang J, Jonkers DMAE, Masclee AAM, Brix S, Li J, Arts I. CW, and Kristiansen K. Impact of early events and lifestyle on the gut microbiota and metabolic phenotypes in young school-age children. Microbiome 2019; 7(1), 1–14. <https://doi.org/10.1186/s40168-018-0608-z>